



[Go to Product page](#)

Datasheet for ABIN2812490

## anti-DOK6 antibody (AA 1-100) (AbBy Fluor® 594)

### Overview

Quantity:	100 µL
Target:	DOK6
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DOK6 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DOK6
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	DOK6
Alternative Name:	DOK6 ( <a href="#">DOK6 Products</a> )
Background:	Synonyms: Docking protein 5, Docking protein 5 like, Docking protein 6, DOK 6, DOK6, DOK-6,

## Target Details

---

DOK5, DOK 5, DOK-5, DOK5L, Downstream of tyrosine kinase 6, HsT3226, MGC20785, Protein dok 5, DOK6\_HUMAN.

Background: The downstream of kinase family (Dok-1-7) are members of a class of docking? proteins that include the tyrosine kinase substrates IRS-1 and Cas, which contain multiple tyrosine residues and putative SH2 binding sites. Dok-4, Dok-5 and Dok-6 are more similar to each other than to the other Dok family members, and may constitute a subfamily of the DOK genes. Dok-5 is a tyrosine kinase substrate that enhances c-Ret-dependent activation of mitogen-activated protein kinase (MAPK). Dok-5 transcript is abundant in muscle and increases during T cell activation. Dok-5 protein undergoes tyrosine phosphorylation in response to insulin and insulin-like growth factor-1. Dok-6 is highly expressed in the developing central nervous system. It associates with Ret to transduce Ret-mediated processes such as axonal projection.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months