antibodies -online.com





anti-DOK7 antibody (AA 21-120) (Alexa Fluor 594)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	DOK7	
Binding Specificity:	AA 21-120	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DOK7 antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DOK7
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	DOK7
Alternative Name:	DOK7 (DOK7 Products)

Target Details

Expiry Date:

-		
Background:	Synonyms: Docking protein 7, DOK 7, DOK7, DOK7_HUMAN, Downstream of tyrosine kinase 7,	
	Protein Dok-7.	
	Background: The downstream of kinase family (Dok1-7) are members of a class of docking?	
	proteins that include the tyrosine kinase substrates IRS-1 and Cas, which contain multiple	
	tyrosine residues and putative SH2 binding sites. Based on their similarities, the Dok family of	
	proteins can be divided into three subgroups: Dok-1/2/3, Dok-4/5/6 and Dok-7. Through its	
	interaction with muscle-specific receptor kinase (MuSK), Dok-7 is crucial for neuromuscular	
	synaptogenesis and for MuSK activation. Mice lacking Dok-7 do not form neuromuscular	
	synapses nor acetylcholine receptor clusters. Mutations in the Dok-7 gene can cause	
	congenital myasthenic syndromes (CMA) recessively inherited disorders characterized by	
	muscle weakness.	
Pathways:	Skeletal Muscle Fiber Development	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Funime Datas	10	

12 months