

Datasheet for ABIN2812549

anti-DUSP10 antibody (AA 161-260) (AbBy Fluor® 594)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DUSP10
Binding Specificity:	AA 161-260
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DUSP10 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DUSP10/MKP5
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DUSP10
Alternative Name:	DUSP10/MKP5 (DUSP10 Products)

Target Details

Background: Synonyms: Dual specificity phosphatase 10, Dual specificity phosphatase MKP 5, Dual specificity phosphatase MKP5, Dual specificity protein phosphatase 10, DUS10_HUMAN, DUSP 10, DUSP10, DUSP10 protein, MAP kinase phosphatase 5, Mitogen activated protein kinase phosphatase 5, Mitogen-activated protein kinase phosphatase 5, MKP 5, MKP-5, MKP5, Serine/threonine specific protein phosphatase.

Background: Mitogen-activated protein (MAP) kinases are a large class of proteins involved in signal transduction pathways that are activated by a range of stimuli and mediate a number of physiological and pathological changes in the cell. Dual specificity phosphatases (DSPs) are a subclass of the protein tyrosine phosphatase (PTP) gene superfamily, which are selective for dephosphorylating critical phosphothreonine and phosphotyrosine residues within MAP kinases. DSP gene expression is induced by a host of growth factors and/or cellular stresses, thereby negatively regulating MAP kinase superfamily members including MAPK/ERK, SAPK/JNK and p38. MKP-5 preferentially binds to p38, but also to SAPK/JNK. It is ubiquitously expressed and localizes to both the cytoplasm and the nucleus. MKP-5 has been implicated in cell proliferation and apoptosis, tumor invasion and immune responses.

Gene ID: 11221

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months