

Datasheet for ABIN2812611

anti-UBR1 antibody (AA 801-900) (AbBy Fluor® 594)



Go to Product page

Overview

Quantity:	100 μL
Target:	UBR1
Binding Specificity:	AA 801-900
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBR1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human eIF3A
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UBR1
Alternative Name:	e3A (UBR1 Products)

Target Details

Background:	Synonyms: EIF3, P167, p180, p185, TIF32, EIF3S10, eIF3-p170, eIF3-theta, Eukaryotic
	translation initiation factor 3 subunit A, eIF3a, Eukaryotic translation initiation factor 3 subunit
	10, eIF-3-theta, eIF3 p167, eIF3 p180, eIF3 p185, KIAA0139
	Background: Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which
	is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates
	with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-
	tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates
	mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3
	complex is also required for disassembly and recycling of post-termination ribosomal
	complexes and subsequently prevents premature joining of the 40S and 60S ribosomal
	subunits prior to initiation.
Gene ID:	8661
UniProt:	Q14152
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months