# antibodies -online.com





## anti-SLC9A3R1 antibody (AA 281-358) (AbBy Fluor® 594)



Go to Product page

/ N	1	-	M	/ 1	01	A /
0	1//	$\vdash$	1 1	/ I	-	<b>\/\/</b>
$\sim$	v	$\sim$	ı ۱	/ I	$\sim$	v v

Quantity:	100 μL
Target:	SLC9A3R1
Binding Specificity:	AA 281-358
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC9A3R1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human EBP50/SLC9A3R1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target:	SLC9A3R1
Alternative Name:	EBP50 (SLC9A3R1 Products)
Background:	Synonyms: EBP 50, EBP50, Ezrin radixin moesin binding phosphoprotein 50, Ezrin-radixin-

moesin-binding phosphoprotein 50, Na+/H+ exchange regulatory cofactor NHE RF, Na+/H+ exchange regulatory cofactor NHE-RF1, Na+/H+ exchange regulatory co factor, NHERF 1, NHERF, NHERF-1, NHERF1, NHRF1\_HUMAN, NPHLOP2, Regulatory cofactor of Na+/H+ exchanger, SLC9A3R1, Sodium hydrogen exchanger regulatory factor 1, Sodium-hydrogen exchanger regulatory factor 1, Solute carrier family 9 sodium/hydrogen exchanger member 3 regulator 1, Solute carrier family 9 sodium/hydrogen exchanger, isoform 3 regulatory factor 1, Solute carrier family 9 isoform 3 regulatory factor 1, Solute carrier family 9 isoform 3 regulatory factor 1, Solute carrier family 9 member 3 regulator 1.

Background: Scaffold protein that connects plasma membrane proteins with members of the ezrin/moesin/radixin family and thereby helps to link them to the actin cytoskeleton and to regulate their surface expression. Necessary for recycling of internalized ADRB2. Was first known to play a role in the regulation of the activity and subcellular location of SLC9A3. Necessary for cAMP-mediated phosphorylation and inhibition of SLC9A3. May enhance Wnt signaling. May participate in HTR4 targeting to microvilli (By similarity). Interacts with MCC.

Gene ID: 9368

Pathways: Proton Transport, Platelet-derived growth Factor Receptor Signaling, Negative Regulation of

Transporter Activity, SARS-CoV-2 Protein Interactome

#### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

### Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months