antibodies.

## Datasheet for ABIN2812669 anti-EDRF1 antibody (AbBy Fluor® 594)



Overview	
Quantity:	100 µL
Target:	EDRF1 (C10orf137)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EDRF1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human EDRF1/C10orf137
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	EDRF1 (C10orf137)
Alternative Name:	EDRF1 (C10orf137 Products)
Background:	Synonyms: C10orf137 protein, Chromosome 10 open reading frame 137, EDRF1, Erythroid
	dferentiation related factor 1, EDRF1_HUMAN.
	Background: The tetratricopeptide repeat (TPR) motif is a degenerate, 34 amino acid sequence
	found in many proteins and acts to mediate protein-protein interactions in various pathways. At

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2812669 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

	the sequence level, there can be up to 16 tandem TPR repeats, each of which has a helix-turn-
	helix shape that stacks on other TPR repeats to achieve ligand binding specificity. EDRF1
	(erythroid differentiation-related factor 1), also known as C10orf137 (chromosome 10 open
	reading frame 137), is a 1,238 amino acid protein containing two TPR repeats. Localizing to
	nucleus, EDRF1 is involved in transcriptional activation of globin genes by regulating DNA-
	binding activity of GATA-1 transcription factor. EDRF1 may also play an important role in organ
	development and histological differentiation. EDRF1 exists as four alternatively spliced
	isoforms and is encoded by a gene mapping to human chromosome 10q26.13.
Gene ID:	26098
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
_	

 Storage:
 -20 °C

 Storage Comment:
 Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date:

12 months