

Datasheet for ABIN2812683

anti-FBLN4 antibody (AA 331-443) (AbBy Fluor® 594)



Go to Product page

Overview	
Quantity:	100 μL
Target:	FBLN4
Binding Specificity:	AA 331-443
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBLN4 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human EFEMP2/Fibulin 4
	KLH conjugated synthetic peptide derived from human EFEMP2/Fibulin 4
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Predicted Reactivity:	IgG Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse
Immunogen: Isotype: Predicted Reactivity: Purification:	IgG Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse
Immunogen: Isotype: Predicted Reactivity: Purification: Target Details	IgG Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse Purified by Protein A.

fibulin-like extracellular matrix protein 2, FBLN 4, FBLN4, FBLN4_HUMAN, FIBL 4, FIBL-4, FIBL4, Fibulin4, Fibulin-4, MBP 1, MBP1, Mutant p53 binding protein 1, Protein UPH1, UPH 1, UPH1, UPH1 protein.

Background: Fibulin proteins contribute to normal development of elastic fiber systems in various types of organs that require elasticity, such as vasculature, lung and skin. Fibulin-4, also known as EFEMP2 (EGF-containing fibulin-like extracellular matrix protein 2), MBP1 or UPH1 is a 443 amino acid secreted protein that contains six EGF-like calcium-binding domains and belongs to the fibulin family. Expressed ubiquitously with highest expression in heart, Fibulin-4 is essential for connective tissue development and elastic fiber formation, and may also play an important role in vascular patterning and collagen biosynthesis. Defects in the gene encoding Fibulin-4 are associated with autosomal recessive cutis laxa type I (CL type I), a connective tissue disorder that is inherited in both an autosomal dominant and an autosomal recessive manner and is characterized by inelastic tissue in all affected areas of the body.

Gene ID:

30008

Application Details

Application Notes:	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months