

Datasheet for ABIN2813043

anti-FAK antibody (pSer910) (AbBy Fluor® 594)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FAK (PTK2)
Binding Specificity:	pSer910
Reactivity:	Mouse, Rat, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAK antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human PTK2 around the phosphorylation site of Tyr910
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat, Xenopus laevis
Predicted Reactivity:	Human,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	FAK (PTK2)
---------	------------

Target Details

Alternative Name:	FAK (PTK2 Products)
Background:	Synonyms: FAK phospho S910, p-FAK phospho S910, FADK 1, FADK, FAK 1, FAK related non kinase polypeptide, FAK1, Focal adhesion kinase 1, FRNK, pp125FAK, Protein tyrosine kinase 2, Protein Tyrosine Kinase Cytoplasmic, PTK 2, FAK1_HUMAN, Focal adhesion kinase-related nonkinase, Protein phosphatase 1 regulatory subunit 71, PPP1R71, Protein-tyrosine kinase 2, p125FAK. Background: Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Microtubule-induced dephosphorylation at Tyr-397 is crucial for the induction of focal adhesion disassembly. Plays a potential role in oncogenic transformations resulting in increased kinase activity.
Gene ID:	5747
Pathways:	Response to Growth Hormone Stimulus , CXCR4-mediated Signaling Events , Smooth Muscle Cell Migration , Signaling of Hepatocyte Growth Factor Receptor , VEGF Signaling

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months