

## Datasheet for ABIN2813052

## anti-FAK antibody (pSer722) (AbBy Fluor® 594)



Go to Product page

()	ve	r\/i	Δ	۱۸/
$\circ$	V C	1 V		v v

Quantity:	100 μL
Target:	FAK (PTK2)
Binding Specificity:	pSer722
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAK antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FAK around the phosphorylation site of Ser722
Isotype:	IgG
Specificity:	Recognizes Ser760 in mice
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	FAK (PTK2)	
Alternative Name:	FAK (PTK2 Products)	
Background:	Synonyms: FAK, FADK, FAK1, FRNK, PPP1R71, p125FAK, pp125FAK, Focal adhesion kinase 1,	
	FADK 1, Focal adhesion kinase-related nonkinase, Protein phosphatase 1 regulatory subunit 71,	
	Protein-tyrosine kinase 2, PTK2	
	Background: This gene encodes a cytoplasmic protein tyrosine kinase which is found	
	concentrated in the focal adhesions that form between cells growing in the presence of	
	extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of	
	protein tyrosine kinases but lacks significant sequence similarity to kinases from other	
	subfamilies. Activation of this gene may be an important early step in cell growth and	
	intracellular signal transduction pathways triggered in response to certain neural peptides or to	
	cell interactions with the extracellular matrix. Several transcript variants encoding different	
	isoforms have been found for this gene, but the full-length natures of only three of them have	
	been determined. [provided by RefSeq, Dec 2010]	
Gene ID:	5747	
UniProt:	Q05397	
Pathways:	Response to Growth Hormone Stimulus, CXCR4-mediated Signaling Events, Smooth Muscle	
	Cell Migration, Signaling of Hepatocyte Growth Factor Receptor, VEGF Signaling	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
	ProClin	

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months