# antibodies -online.com





## anti-FBXW12 antibody (Alexa Fluor 594)



( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	FBXW12	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FBXW12 antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FBXW12/FBX035	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	FBXW12	
Alternative Name:	FBXW12 (FBXW12 Products)	
Background:	Synonyms: F box and WD 40 domain containing protein 12; F box and WD 40 domain protein	
	12; F box and WD repeat domain containing 12; F box and WD40 repeat containing protein; F	
	box only protein 35; F box/WD repeat containing protein 12; F-box and WD-40 domain-	
	containing protein 12; F-box only protein 35; F-box/WD repeat-containing protein 12; Fbw12;	

FBW12\_HUMAN; FBX12; FBXO12; FBXO35; FBXW 12; FBXW12.

Background: Belonging to the F-box family of proteins, FBXO35 (F-box only protein 35), also designated F-box and WD-40 domain-containing protein 12 (FBXW12 or FBXO12), is a 464 amino acid protein that contains one C-terminal F-box domain. F-box proteins are critical components of the SCF (Skp1-CUL-1-F-box protein)-type E3 ubiquitin ligase complex and are involved in substrate recognition and recruitment for ubiquitination. They are members of a larger family of proteins that are involved in the regulation of a wide variety of cellular mechanisms, including the cell cycle, the immune response, signaling cascades and developmental processes. They function by targeting proteins, such as cyclins, cyclindependent kinase inhibitors, I°B-å and -catenin, for degradation by the proteasome after ubiquitination. Via its F-box domain, FBXO35 can directly interact with Skp1 p19 and CUL-1. FBXO35 is ubiquitously expressed at low levels in most human tissues.

Gene ID:

285231

IF(IHC-P) 1:50-200

#### **Application Details**

Application Notes:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months