antibodies .- online.com







anti-GADD34 antibody (AA 401-500) (AbBy Fluor® 594)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	GADD34 (PPP1R15A)	
Binding Specificity:	AA 401-500	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GADD34 antibody is conjugated to AbBy Fluor® 594	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GADD34	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	GADD34 (PPP1R15A)
Alternative Name:	GADD34 (PPP1R15A Products)
Background: Synonyms: GADD34, Protein phosphatase 1 regulatory subunit 15A, Growth arrest and [

Expiry Date:

Target Details	
	damage-inducible protein GADD34, Myeloid differentiation primary response protein MyD116 homolog, PPP1R15A
	Background: Recruits the serine/threonine-protein phosphatase PP1 to dephosphorylate the translation initiation factor eIF-2A/EIF2S1, thereby reversing the shut-off of protein synthesis
	initiated by stress-inducible kinases and facilitating recovery of cells from stress. Down-regulates the TGF-beta signaling pathway by promoting dephosphorylation of TGFB1 by PP1. May promote apoptosis by inducing TP53 phosphorylation on 'Ser-15'.
Gene ID:	23645
UniProt:	075807
Pathways:	ER-Nucleus Signaling, Unfolded Protein Response
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months