

Datasheet for ABIN2813573

anti-GABRB1 antibody (pSer434) (AbBy Fluor® 594)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	GABRB1
Binding Specificity:	pSer434
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GABRB1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human GARB1 around the phosphorylation site of Ser434
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GABRB1
---------	--------

Target Details

Alternative Name:	GARB1 (GABRB1 Products)
Background:	Synonyms: GABA A Receptor beta 1 phospho S434; GARB1 Ser 434; GABAA receptor subunit beta-1; GABA-A receptor, beta-1 polypeptide; Gabrb-1; Gamma Aminobutyric Acid A Receptor Beta 1; Gamma Aminobutyric Acid Receptor , beta-1; Gamma-aminobutyric acid GABA A receptor, subunit beta 1; Gamma-aminobutyric acid receptor subunit beta-1; GARB1; GABRA1; AW061132; B230208N19Rik; GABAA receptor beta 1; GABAA receptor subunit beta-1; GABA-A receptor, beta-1 polypeptide; Gabrb-1; GABRB1; Gamma aminobutyric acid GABA A receptor beta 1; Gamma Aminobutyric Acid A Receptor Beta 1; Gamma Aminobutyric Acid Receptor , beta-1; Gamma-aminobutyric acid GABA A receptor, subunit beta 1; Gamma-aminobutyric acid receptor subunit beta-1; GARB1; GBRB1_HUMAN. Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (g-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl⁻ conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R alpha 1-6, GABAA R beta 1-3, GABAA Rρ1-3, GABAA Rρ, GABAA R gamma, GABAA R delta 1 and GABAA R delta 2. The GABAB family is composed of GABAB R1 alpha and GABAB R1 beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2 and -3). The GABA transporters function to terminate GABA action.

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months