



[Go to Product page](#)

Datasheet for ABIN2813681

anti-GEM antibody (AA 201-296) (AbBy Fluor® 594)

Overview

Quantity:	100 µL
Target:	GEM
Binding Specificity:	AA 201-296
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GEM antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GEM
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	GEM
Alternative Name:	GEM (GEM Products)

Target Details

Background: Synonyms: GTP binding mitogen induced T cell protein, GTP binding protein expressed in mitogen stimulated T cells, GTP binding protein GEM, GTP binding protein overexpressed in skeletal muscle, Kinase inducible Ras like protein, KIR, MGC26294, RAS like protein KIR, GEM_HUMAN.

Background: Gem belongs to the Rad/Gem/Kir (RGK) subfamily of Ras-related GTPases, which lack typical C-terminal amino acid motifs for isoprenylation. Rad and Gem bind calmodulin in a Ca²⁺-dependent manner via this C-terminal extension, involving residues 278-297 in human Rad. High intracellular Gem levels, which interact with intact microtubules and microfilaments, promote profound changes in cell morphology. Ectopic Gem expression is sufficient to stimulate cell flattening and neurite extension in N1E-115 and SH-SY5Y neuroblastoma cells, suggesting a role for Gem in cytoskeletal rearrangement and/or morphological differentiation of neurons. Gem was also observed in developing trigeminal nerve ganglia in 12.5 day mouse embryos, demonstrating that Gem expression is a property of normal ganglionic development. The interaction of Gem with beta-subunits regulates Ca²⁺ channel expression at the cell surface. The human Gem gene maps to chromosome ,8q22.1.

Gene ID: 2669

Application Details

Application Notes: FCM 1:20-100
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months