antibodies .- online.com





anti-KCNJ9 antibody (AA 61-160) (Alexa Fluor 594)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	KCNJ9
Binding Specificity:	AA 61-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNJ9 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GIRK3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target:	KCNJ9
Alternative Name:	GIRK3 (KCNJ9 Products)
Background: Synonyms: G protein activated inward rectier potassium channel 3, G protein coupled inwa	

rectier potassium channel, GIRK3, Inwardly rectier K+ channel Kir3.3, Inwardly rectier K+ channel KIR3.3, KIR3.3, Potassium channel inwardly rectying subfamily J member 9, Potassium inwardly rectying channel subfamily J member 9, Potassium inwardly rectying channel subfamily J9.

Background: KCNJ9 belongs to the inward rectifier-type potassium channel family and is controlled by G proteins. It associates with another G-protein-activated potassium channel to form a heteromultimeric pore-forming complex. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium, as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium.

Gene ID:

3760

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months