antibodies -online.com





anti-HIRA antibody (AA 251-350) (AbBy Fluor® 594)



0	V	el	V	İΕ	۷۶	١

Quantity:	100 μL	
Target:	HIRA	
Binding Specificity:	AA 251-350	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HIRA antibody is conjugated to AbBy Fluor® 594	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HIRA/DGGR1	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	HIRA
Alternative Name:	HIRA (HIRA Products)

Target Details

Background:	Synonyms: DGCR1, DGGR 1, DGGR1, DiGeorge critical region gene 1, HIR, HIR histone cell cycle
	regulation defective homolog A, HIRA, HIRA protein, HIRA_HUMAN, Protein HIRA, TUP 1, TUP1,
	TUP1 like enhancer of split protein 1, TUP1-like enhancer of split protein 1, TUPLE 1, TUPLE1.
	Background: The HIRA gene encodes a histone chaperone that preferentially places the variant
	histone H3.3 in nucleosomes. Orthologs of this gene in flies, yeast and plants are necessary for
	the formation of transcriptionally silent heterochomatin. This gene plays an important role in
	the formation of the senescence-associated heterochromatin foci. These foci likely mediate the
	irreversible cell cycle changes that occur in senescent cells. It is considered the primary
	candidate gene in some haploinsufficiency syndromes such as DiGeorge syndrome, and
	insufficient production of the gene may disrupt normal embryonic development.
Gene ID:	7290
Pathways:	Chromatin Binding
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months