

Datasheet for ABIN2814627

anti-IDH3B antibody (AA 151-250) (AbBy Fluor® 594)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IDH3B
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IDH3B antibody is conjugated to AbBy Fluor® 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IDH3B
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	IDH3B
Alternative Name:	IDH3B (IDH3B Products)
Background:	Synonyms: FLJ11043, H-IDHB, IDH3B, IDH3B_HUMAN, Isocitrate dehydrogenase [NAD] subunit

Target Details

beta, Isocitrate dehydrogenase [NAD] subunit beta, mitochondrial, isocitrate dehydrogenase 3, beta subunit, Isocitric dehydrogenase, Isocitric dehydrogenase subunit beta, MGC903, mitochondrial, NAD+-specic ICDH, NAD+-specic ICDH subunit beta, NAD+-specic isocitrate dehydrogenase b subunit, NAD+-specic isocitrate dehydrogenase beta, OTTHUMP00000030023_ OTTHUMP00000030024, RP46.

Background: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. The protein encoded by this gene is the beta subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase. Three alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008].

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months