

Datasheet for ABIN2814694

anti-IKZF1 antibody (AA 251-360) (AbBy Fluor® 594)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IKZF1
Binding Specificity:	AA 251-360
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IKZF1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IKAROS
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	IKZF1
Alternative Name:	IKAROS (IKZF1 Products)

Target Details

Background:	Synonyms: IKAROS family zinc finger 1 Ikaros, PRO0758, CLL associated antigen KW 6, DNA-binding protein Ikaros, hlk 1, hlk-1, IK1, Ikaros zinc finger protein, IKAROS, Ikaros family zinc finger protein 1, Ikzf1, IKZF1_HUMAN, LYF1, Lymphoid transcription factor LyF-1, zinc finger protein subfamily 1A 1, ZNFN1A1. Background: Transcription regulator of hematopoietic cell differentiation. Binds gamma-satellite DNA. Binds with higher affinity to gamma satellite A. Plays a role in the development of lymphocytes, B- and T-cells. Binds and activates the enhancer (delta-A element) of the CD3-delta gene. Repressor of the TDT (terminal deoxynucleotidyltransferase) gene during thymocyte differentiation. Regulates transcription through association with both HDAC-dependent and HDAC-independent complexes. Targets the 2 chromatin-remodeling complexes, NuRD and BAF (SWI/SNF), in a single complex (PYR complex), to the beta-globin locus in adult erythrocytes. Increases normal apoptosis in adult erythroid cells. Confers early temporal competence to retinal progenitor cells (RPCs).Tissue specificity:Abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. Lower expression in bone marrow and small intestine.
-------------	---

Gene ID:	10320
----------	-------

Pathways:	Production of Molecular Mediator of Immune Response
-----------	---

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months