
antibodies -online.com





anti-CYFIP2 antibody






Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Overview		
Quantity:	100 μL	
Target:	CYFIP2	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CYFIP2 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the center region of human CYFIP2.	
	The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Characteristics:	Rabbit Polyclonal antibody to CYFIP2 (cytoplasmic FMR1 interacting protein 2)	
	CYFIP2 antibody [N2C1], Internal	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	CYFIP2	
Alternative Name:	cytoplasmic FMR1 interacting protein 2 (CYFIP2 Products)	

Target Details

Dooleground	Collular Lacelization: Ottoploom parinusloom reside Collingation are served.	
Background:	Cellular Localization: Cytoplasm , perinuclear region , Cell junction , synapse , synaptosome	
Molecular Weight:	148 kDa	
Gene ID:	26999	
UniProt:	Q96F07	
Pathways:	RTK Signaling, Apoptosis, Positive Regulation of Endopeptidase Activity	
Application Details		
Application Notes:	WB: 1:5000-1:20000. Optimal dilutions/concentrations should be determined by the researche Not tested in other applications.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1.5 mg/mL	
Buffer:	1XPBS pH 7, 20 % Glycerol, 0.025 % ProClin 300	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Publications		
Product cited in:	Rodriguez-Gil, Hu, Greer: "Dishevelled proteins are associated with olfactory sensory neuron presynaptic terminals." in: PLoS ONE , Vol. 8, Issue 2, pp. e56561, (2013) (PubMed).	



Western Blotting

Image 1. WB Image CYFIP2 antibody detects CYFIP2 protein by western blot analysis. Various whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with CYFIP2 antibody , diluted at a dilution of 1:10000.