

Datasheet for ABIN2854616
anti-PRKG1 antibody (Center)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	PRKG1
Binding Specificity:	Center
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRKG1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Recombinant protein encompassing a sequence within the center region of human cGK1. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human, Cow (Bovine)
Cross-Reactivity (Details):	Human (100 %), Bovine (100 %)
Characteristics:	Rabbit Polyclonal antibody to cGK1 (protein kinase, cGMP-dependent, type I) cGK1 antibody
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	PRKG1
Alternative Name:	cGK1 (PRKG1 Products)
Background:	<p>Mammals have three different isoforms of cyclic GMP-dependent protein kinase (Ialpha, Ibeta, and II). These PRKG isoforms act as key mediators of the nitric oxide/cGMP signaling pathway and are important components of many signal transduction processes in diverse cell types.</p> <p>This PRKG1 gene on human chromosome 10 encodes the soluble Ialpha and Ibeta isoforms of PRKG by alternative transcript splicing. A separate gene on human chromosome 4, PRKG2, encodes the membrane-bound PRKG isoform II. The PRKG1 proteins play a central role in regulating cardiovascular and neuronal functions in addition to relaxing smooth muscle tone, preventing platelet aggregation, and modulating cell growth. This gene is most strongly expressed in all types of smooth muscle, platelets, cerebellar Purkinje cells, hippocampal neurons, and the lateral amygdala. Isoforms Ialpha and Ibeta have identical cGMP-binding and catalytic domains but differ in their leucine/isoleucine zipper and autoinhibitory sequences and therefore differ in their dimerization substrates and kinase enzyme activity. [provided by RefSeq, Sep 2011]</p>
Molecular Weight:	76 kDa
Gene ID:	5592

Application Details

Application Notes:	<p>Suggested dilution Reference Western blot 1:5000-1:20000* Not tested in other applications.</p> <p>*Optimal dilutions/concentrations should be determined by the researcher.Suggested dilutionReferenceWestern blot1:5000-1:20000*</p>
Comment:	Positive Control: JC , BCL-1
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.1M Tris, 0.1M Glycine, 10 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative.
Preservative:	Thimerosal (Merthiolate)

Handling

Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Validation report #104404 for Cleavage Under Targets and Release Using Nuclease (CUT&RUN)



Western Blotting

Image 1. WB Image Sample (30 ug of whole cell lysate) A: JC B: BCL-1 7.5% SDS PAGE antibody diluted at 1:10000