# ANTIBODIES ONLINE

Datasheet for ABIN2855042 anti-CTNNB1 antibody (N-Term)				
1 Validation	14 Images	8 Publications		
Overview			do to riodact page	
Quantity:	100 µL			
Target:	CTNNB1			
Binding Specificity:	N-Term			
Reactivity:	Human			
Host:	Rabbit			
Clonality:	Polyclonal			
Conjugate:	This CTNNB1 antibody is un-con	jugated		
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro)), Chromatin Immunoprecipitation (ChIP), Immunohistochemistry (Whole Mount) (IHC (wm)), Cleavage Under Targets and Release Using Nuclease (CUT&RUN)			

# Product Details

Immunogen:	Recombinant protein encompassing a sequence within the N-terminus region of human beta Catenin. The exact sequence is proprietary.
Isotype:	lgG
Cross-Reactivity:	Cat, Dog, Human, Mouse, Rabbit, Rat, Zebrafish (Danio rerio)
Characteristics:	Rabbit Polyclonal antibody to beta Catenin (catenin (cadherin-associated protein), beta 1, 88 kDa) beta Catenin antibody [N1N2-2], N-term

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/9 | Product datasheet for ABIN2855042 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Product Details	
Purification:	Purified by antigen-affinity chromatography.
Grade:	KO Validated
Target Details	
Target:	CTNNB1
Alternative Name:	catenin beta 1 (CTNNB1 Products)
Background:	Beta-catenin is an adherens junction protein. Adherens junctions (AJs, also called the zonula adherens) are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces. AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior. At several stages of embryogenesis, wound healing, and tumor cell metastasis, cells form and leave epithelia. This process, which involves the disruption and reestablishment of epithelial cell-cell contacts, may be regulated by the disassembly and assembly of AJs. AJs may also function in the transmission of the 'contact inhibition' signal, which instructs cells to stop dividing once an epithelial sheet is complete.[supplied by OMIM]
Molecular Weight:	85 kDa
Gene ID:	1499
UniProt:	P35222
Pathways:	WNT Signaling, Intracellular Steroid Hormone Receptor Signaling Pathway, Peptide Hormone Metabolism, Regulation of Muscle Cell Differentiation, Cell-Cell Junction Organization, Tube Formation, Maintenance of Protein Location, Signaling Events mediated by VEGFR1 and VEGFR2
Application Details	
Application Notes:	WB: 1:500-1:20000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. FACS: 1:50-1:200. IP: 1:50-1:100. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: Mouse brain , PC-12 , HeLa Validation: Comparison, KO/KD, Orthogonal

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/9 | Product datasheet for ABIN2855042 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

## Application Details

#### Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	0.07 mg/mL
Buffer:	1XPBS ( pH 7), 1 % BSA, 20 % Glycerol, 0.025 % ProClin 300
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

## Publications

Product cited in:Noack, Bundkirchen, Xu, Gylstorff, Zhou, Köhler, Jantaree, Neunaber, Nowak, Relja: "AcuteIntoxication With Alcohol Reduces Trauma-Induced Proinflammatory Response and BarrierBreakdown in the Lung via the Wnt/β-Catenin Signaling Pathway." in: Frontiers in immunology,Vol. 13, pp. 866925, (2022) (PubMed).

Zambanini, Nordin, Jonasson, Pagella, Cantù: "A new cut&run low volume-urea (LoV-U) protocol optimized for transcriptional co-factors uncovers Wnt/b-catenin tissue-specific genomic targets." in: **Development (Cambridge, England)**, (2022) (PubMed).

Yin, Yao, Deng, Wang, Cai, Shen: "Identification of candidate IncRNAs and circRNAs regulating WNT3/β-catenin signaling in essential hypertension." in: **Aging**, Vol. 12, Issue 9, pp. 8261-8288, (2020) (PubMed).

Fang, Lin, Liang, Liang: "A novel c-Kit/phospho-prohibitin axis enhances ovarian cancer stemness and chemoresistance via Notch3-PBX1 and β-catenin-ABCG2 signaling." in: **Journal of biomedical science**, Vol. 27, Issue 1, pp. 42, (2020) (PubMed).

Lin, Lee, Wu, Wu, Chen, Huang, Chang, Cheng, Chang, Ho, Tu, Ho: "Small G protein signalling

modulator 2 (SGSM2) is involved in oestrogen receptor-positive breast cancer metastasis through enhancement of migratory cell adhesion via interaction with E-cadherin." in: **Cell adhesion & migration**, Vol. 13, Issue 1, pp. 120-137, (2019) (PubMed).

There are more publications referencing this product on: Product page

#### Images



#### Western Blotting

**Image 1.** WB Image beta Catenin antibody [N1N2-2], N-term detects beta Catenin protein by western blot analysis. Various whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with beta Catenin antibody [N1N2-2], N-term , diluted at 1:1000.

#### Western Blotting

Image 2. WB Image Sample (30 ug of whole cell lysate) A: HeLa 7.5% SDS PAGE antibody diluted at 1:1000

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/9 | Product datasheet for ABIN2855042 | 07/26/2024 | Copyright antibodies-online. All rights reserved.



#### Immunohistochemistry

**Image 3.** IHC-P Image beta Catenin antibody [N1N2-2], Nterm detects beta Catenin protein at membrane on mouse urinary bladder by immunohistochemical analysis. Sample: Paraffin-embedded mouse urinary bladder. beta Catenin antibody [N1N2-2], N-term , diluted at 1:500.

Please check the product details page for more images. Overall 14 images are available for ABIN2855042.

Validation report #104235 for Cleavage Under Targets and Release Using Nuclease (CUT&RUN)



## Successfully validated (Cleavage Under Targets and Release Using Nuclease (CUT&RUN))

by Cantù Lab, Gene Regulation during Development and Disease, Linköping University Report Number: 104235 Date: Jun 28 2021

Target:	beta Catenin	
Lot Number:	42536	
Method validated:	Cleavage Under Targets and Release Using Nuclease (CUT&RUN)	
Positive Control:	Recombinant anti-H3K27me3 CUT&RUN Positive Control antibody (antibodies-online, ABIN6923144)	
Negative Control:	Monoclonal anti-FLAG (Sigma-Aldrich, F3165)	
Notes:	Passed. ABIN2855042 allows for beta Catenin-targeted digestion using CUT&RUN on HEK-293 nuclei.	
Primary Antibody:	ABIN2855042	
Protocol:	<ul> <li>Cell harvest</li> <li>Harvest 500,000 HEK293T cells per antibody to be used at RT.</li> <li>Centrifuge cell solution 3 min at 600 x g at RT.</li> <li>Remove the liquid carefully.</li> <li>Gently resuspend cells in 1 mL Wash Buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 0.5 mM Spermidine, Roche Complete Protease Inhibitor EDTA-free) by pipetting and transfer cell solution to a 2 mL microcentrifuge tube.</li> <li>Centrifuge cell solution 3 min at 600 x g at RT and discard the supernatant.</li> <li>Repeat twice for a total of three washes.</li> <li>Resuspend cell pellet in 1 mL Wash Buffer by gently pipetting.</li> <li>Concanavalin A beads preparation</li> <li>Prepare one 1.5 mL microcentrifuge tube.</li> <li>Gently resuspend the magnetic Concanavalin A Beads (antibodies-online, ABIN6923139).</li> <li>Pipette 40 µL Con A Beads slurry for each sample into the 1.5 mL microcentrifuge tube.</li> <li>Place the tube on a magnet stand until the fluid is clear. Remove the liquid carefully.</li> <li>Remove the microcentrifuge tube from the magnetic stand.</li> <li>Pipette 1 mL Binding Buffer (20 mM HEPES pH 7.5, 10 mM KCl, 1 mM CaCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>) into each tube and resuspend ConA beads by gentle pipetting.</li> <li>Spin down the liquid from the lid with a quick pulse in a table-top centrifuge.</li> <li>Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.</li> <li>Remove the microcentrifuge tube from the magnetic stand.</li> </ul>	

- Repeat twice for a total of three washes.
- Gently resuspend the ConA Beads in a volume of Binding Buffer corresponding to the original volume of bead slurry, i.e. 40 µL per sample.
- Nuclei immobilization binding to Concanavalin A beads
  - $\circ~$  Carefully vortex the cell suspension and add 10  $\mu L$  of the Con A beads in Binding Buffer to the cell suspension for each sample.
  - Close tube tightly and rotate for 10 min at RT.
- Primary antibody binding
  - oDivide nuclei suspension into separate 2 mL microcentrifuge tubes, one for each antibody (500,000 nuclei per sample).
  - Place the microcentrifuge tubes on a magnetic stand until the fluid is clear. Remove the liquid carefully.
  - Remove the microcentrifuge tubes from the magnetic stand.
  - Place each tube at a low angle on the vortex mixer set to a low speed and add 150 µL
     Digitonin Wash buffer (wash buffer with 0.025% (wt/vol) Digitonin) supplemented with 2 mM EDTA.
  - Gently vortex the microcentrifuge tubes until the beads are resuspended.
  - Add 1.5 µL antibody (anti-beta Catenin antibody ABIN2855042, anti-H3K27me3 positive control antibody ABIN6923144, and anti-FLAG tag antibody negative control) to the respective tube, corresponding to a 1:100 dilution.
  - $\circ~$  Rotate the microcentrifuge tubes ON at 4 °C.
  - Spin down the liquid and place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
  - Remove the microcentrifuge tubes from the magnetic stand.
  - Resuspend with 1 mL Digitonin Wash Buffer and mix by inversion. If clumping occurs, gently remove the clumps with a 1 ml pipette tip.
  - Repeat once for a total of two washes.
- pAG-MNase Binding
  - Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
  - Remove the microcentrifuge tubes from the magnetic stand.
  - $\circ~$  o Vortex the sample at low speed and add 2.5  $\mu$ L (0.05 volumes) CUTANA pAG-MNase for ChIC/CUT&RUN Assays (ABIN6950951) in 50  $\mu$ L Digitonin Wash Buffer per sample, gently resuspending the beads by pipetting.
  - Rotate the microcentrifuge tubes for 1 h at 4 °C.
  - Spin down the liquid and place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
  - Remove the microcentrifuge tubes from the magnetic stand.
  - Resuspend with 1 mL Digitonin Wash Buffer and mix by inversion. If clumping occurs, gently remove the clumps with a 1 mL pipette tip.
  - Repeat once for a total of two washes.
- MNase digestion and release of pAG-MNase-antibody-chromatin complexes
- Spin down the liquid from the lid with a quick pulse in a table-top centrifuge.
- Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.

- Place each tube at a low angle on the vortex mixer set to a low speed and add 100 µL
   Digitonin Wash buffer per sample along the side of the tube.
- Place tubes in a heat block, kept on ice, and allow to chill.
- $\circ~$  Add 2  $\mu L$  0.1 M CaCl\_2 to each sample.
- Incubate tubes at 0 °C for 30 min.
- Add 100 µL 2xSTOP buffer (340 mM NaCl, 20 mM EDTA, 4 mM EGTA, 0.05% (wt/vol) Digitonin, 100 µg/mL RNAse A, 50 µg/mL Glycogen).
- $\circ~$  Incubate tubes at 37 °C for 30 min.
- Place the tubes on a magnet stand until the fluid is clear.
- Transfer the supernatant containing the pA-MNase-bound digested chromatin fragments to fresh 1.5 mL microcentrifuge tubes.
- DNA extraction
  - $\circ~$  Add 2  $\mu L$  10% SDS to a final concentration of 0.1% and 2.5  $\mu L$  Proteinase K (20 mg/mL) to each supernatant.
  - Gently vortex tubes at a low speed of approximately 1,100 rpm.
  - $\circ~$  Incubate tubes at 50 °C for 1 h.
  - $\circ~$  Add 200  $\mu L$  PCI to tube.
  - Vortex tubes thoroughly at high speed until the liquid appears milky.
  - $\circ~$  Centrifuge tubes in a tabletop centrifuge at 16,000 x g at RT for 5 min.
  - Carefully transfer to upper aqueous phase to a fresh 1.5 mL microcentrifuge tube containing 2 μL glycogen (diluted 1:10 to 2 mg/mL from the 20 mg/mL stock solution).
  - $\circ~$  Add 20  $\mu L$  3 M NaOAc pH 5.2.
  - $\circ~$  Add 400  $\mu L$  100% ethanol.
  - $\circ~$  Place tubes for at -20 °C ON.
  - Centrifuge tubes in a tabletop centrifuge at 16,000 x g at 4 °C for 5min.
  - Remove the liquid carefully with a pipette.
  - Wash pellet with 1ml 70% ethanol.
  - Centrifuge tubes in a tabletop centrifuge at 16,000 x g at 4 °C for 1 min.
  - Remove the liquid carefully with a pipette.
  - $\circ~$  Air-dry the pellet, then dissolve in 30  $\mu L$  1 mM Tris-HCl, 0.1 mM EDTA.
- Library preparation and sequencing
  - Prepare libraries using KAPA HyperPrep Kit using KAPA Dual-Indexed adapters according to protocol.
  - Sequence libraries on an Illumina NextSeq 500 sequencer, using a NextSeq 500/550 High Output Kit v2.5 (75 Cycles), 36bp PE.
- Peak calling
  - Map reads to the GRCm38 (mm10) mouse genome using Bowtie2 with options: --local -very-sensitive- local --no-unal --no-mixed --no-discordant.
  - Call peaks using MACS2 with options -f BAMPE --keep-dup all -nomodel.

Experimental Notes:

Peaks generated using ABIN2855074 for CUT&RUN aligned well with beta Catenin ChIP-seq tracks(Doumpas et al., 2019).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 8/9 | Product datasheet for ABIN2855042 | 07/26/2024 | Copyright antibodies-online. All rights reserved.



1

Validation image no. 1 for anti-Catenin (Cadherin-Associated Protein), beta 1, 88kDa (CTNNB1) (N-Term) antibody (ABIN2855042)

Bioanalyzer profiles comparing fragment size distributions between reads obtained from CUT&RUN using an anti-H3K27me3 CUT&RUN Positive Control antibody ABIN6923144 and anti-beta Catenin ABIN2855042 after library preparation.



Validation image no. 2 for anti-Catenin (Cadherin-Associated Protein), beta 1, 88kDa (CTNNB1) (N-Term) antibody (ABIN2855042)

Alignment tracks from CUT&RUN targeting beta Catenin in HEK293T cells. 1. Alignment track for CUT&RUN reads obtained using anti-beta Catenin antibody ABIN2855042 in HEK293T cells. 2. Peaks called by SEACR from CUT&RUN data using anti-beta Catenin antibody ABIN2855042. 3. Peak track from ChIP-seq targeting beta Catenin in HEK293T cells (Doumpas et al., 2019). 4. Refseq track showing the AXIN2 gene on (chr17:65528563-65561648).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 9/9 | Product datasheet for ABIN2855042 | 07/26/2024 | Copyright antibodies-online. All rights reserved.