

Datasheet for ABIN2855375  
**anti-STK17A antibody (C-Term)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	100 µL
Target:	STK17A
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STK17A antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	Recombinant protein encompassing a sequence within the C-terminus region of human DRAK1. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Characteristics:	Rabbit polyclonal antibody to DRAK1 (serine/threonine kinase 17a) DRAK1 antibody [C1C3]
Purification:	Purified by antigen-affinity chromatography.

## Target Details

Target:	STK17A
---------	--------

## Target Details


Alternative Name:	serine/threonine kinase 17a ( <a href="#">STK17A Products</a> )
Background:	<p>This gene is a member of the DAP kinase-related apoptosis-inducing protein kinase family and encodes an autophosphorylated nuclear protein with a protein kinase domain. The protein has apoptosis-inducing activity.</p> <p>Cellular Localization: Nucleus</p>
Molecular Weight:	47 kDa
Gene ID:	9263
UniProt:	<a href="#">Q9UEE5</a>

## Application Details

Application Notes:	<p>WB: 1:500-1:3000. Optimal dilutions/concentrations should be determined by the researcher.</p> <p>Not tested in other applications.</p>
Comment:	Positive Control: HepG2
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	0.84 mg/mL
Buffer:	0.1M Tris-Glycine ( pH 7), 10 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.



Western Blotting

**Image 1.** WB Image Sample (30 ug of whole cell lysate) A:  
Hep G2 , 10% SDS PAGE DRAK1 antibody antibody diluted  
at 1:500