

Datasheet for ABIN2855752
anti-LRPAP1 antibody (Center)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	LRPAP1
Binding Specificity:	Center
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRPAP1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Recombinant protein encompassing a sequence within the center region of human RAP. The exact sequence is proprietary.
Isotype:	IgG
Characteristics:	Rabbit Polyclonal antibody to RAP (low density lipoprotein receptor-related protein 1) RAP antibody [N1C3]
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	LRPAP1
Alternative Name:	RAP (LRPAP1 Products)

Target Details

Background: Interacts with LRP1/alpha-2-macroglobulin receptor and glycoprotein 330.

Cellular Localization: Endoplasmic reticulum , Cytoplasm , Cell surface

Molecular Weight: 41 kDa

Gene ID: 4043

Application Details

Application Notes: Suggested dilution Reference Western blot 1:500-1:3000* Not tested in other applications.

Optimal dilutions/concentrations should be determined by the researcher.Suggested dilutionReferenceWestern blot1:500-1:3000

Comment: Positive Control: A549 , HeLa , HepG2 , HCT116

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL


Buffer: 1XPBS, 1 % BSA, 20 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative.

Preservative: Thimerosal (Merthiolate)

Precaution of Use: This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.



Western Blotting

Image 1. WB Image Various whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with RAP antibody [N1C3] , diluted at 1:1000.



Western Blotting

Image 2. WB Image Various tissue extracts (50 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with RAP antibody [N1C3] , diluted at 1:1000.