antibodies .- online.com







anti-IL17RD antibody

Images




\sim	
()\/⊝	rview
0 1 0	VICVV

Quantity:	100 μL
Target:	IL17RD
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL17RD antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the center region of human IL17
	receptor D. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Characteristics:	Rabbit Polyclonal antibody to IL17 receptor D (interleukin 17 receptor D)
	IL17 receptor D antibody [N3C2], Internal
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	IL17RD
Alternative Name:	interleukin 17 receptor D (IL17RD Products)


Target Details

Background:	Fibroblast growth factors (FGFs, see MIM 603726) are secreted proteins involved in cellular
	proliferation, migration, differentiation, and survival. FGF activity is negatively regulated by
	members of the 'sprouty' family (e.g., SPRY1, MIM 602465). The SEF protein is a modulator of
	FGF signaling.[supplied by OMIM]
	Cellular Localization: Golgi apparatus membrane, Single-pass type I membrane protein , Cell
	membrane, Single-pass type I membrane protein , Isoform 4: Cytoplasm
Molecular Weight:	82 kDa
Gene ID:	54756
UniProt:	Q8NFM7
Pathways:	Toll-Like Receptors Cascades
Application Details	
Application Notes:	WB: 1:500-1:3000. Optimal dilutions/concentrations should be determined by the researcher.
	Not tested in other applications.
Comment:	Positive Control: Mouse brain , 293T , NIH-3T3
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.85 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 10 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE
	which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage
	(1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid
	multiple freeze-thaw cycles.



Western Blotting

Image 1. WB Image Sample (30 ug of whole cell lysate) A:NIH-3T3 7.5% SDS PAGE antibody diluted at 1:2000



Western Blotting

Image 2. WB Image Sample(30 μ g of whole cell lysate) A:293T 7.5% SDS PAGE antibody diluted at 1:500

Western Blotting

Image 3. WB Image Sample (50 ug of whole cell lysate) A: Mouse brain 7.5% SDS PAGE antibody diluted at 1:1000