antibodies - online.com







anti-BCL2L12 antibody (C-Term)



Images



Overview	
Quantity:	100 μL
Target:	BCL2L12
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BCL2L12 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
lmmunogen:	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human BCL2L12. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Characteristics:	Rabbit Polyclonal antibody to BCL2L12 (BCL2-like 12 (proline rich)) BCL2L12 antibody
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	BCL2L12

Target Details


Alternative Name:	BCL2 like 12 (BCL2L12 Products)
Molecular Weight:	37 kDa
Gene ID:	83596
UniProt:	Q9HB09

Application Details

Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined
	by the researcher. Not tested in other applications.
Comment:	Positive Control: Molt-4
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.37 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 10 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.




Western Blotting

Image 1. WB Image Sample (30 ug of whole cell lysate) A: Molt-4, 10% SDS PAGE antibody diluted at 1:1000



Immunohistochemistry

Image 2. IHC-P Image BCL2L12 antibody detects BCL2L12 protein at cytosol and nucleus on human colon carcinoma by immunohistochemical analysis. Sample: Paraffinembedded colon carcinoma tissue. BCL2L12 antibody, dilution: 1:500.



Immunohistochemistry

Image 3. IHC-P Image BCL2L12 antibody detects BCL2L12 protein at cytosol on U87 xenograft by immunohistochemical analysis. Sample: Paraffin-embedded U87 xenograft. BCL2L12 antibody, dilution: 1:500.