antibodies -online.com





anti-FZD9 antibody (N-Term)





\sim			
()/	$/\Delta r$	\/I <i>C</i>	۸۱۵


Overview	
Quantity:	100 μL
Target:	FZD9
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FZD9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	Carrier-protein conjugated synthetic peptide encompassing a sequence within the N-terminus region of human Frizzled 9. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Mouse (Murine), Rat (Rattus)
Cross-Reactivity (Details):	Mouse (92 %), Rat (92 %)
Characteristics:	Rabbit Polyclonal antibody to Frizzled 9 (frizzled homolog 9 (Drosophila)) Frizzled 9 antibody [N1N2], N-term
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	FZD9	
Alternative Name:	Frizzled 9 (FZD9 Products)	
Background:	Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD9 gene is located within the Williams syndrome common deletion region of chromosome 7, and heterozygous deletion of the FZD9 gene may contribute to the Williams syndrome phenotype. FZD9 is expressed predominantly in brain, testis, eye, skeletal muscle, and kidney.	
	Cellular Localization: Cell membrane, Multi-pass membrane protein	
Molecular Weight:	64 kDa	
Gene ID:	8326	
Pathways:	WNT Signaling	
Application Details		
Application Notes:	Suggested dilution Reference IHC (Formalin-fixed paraffin-embedded sections) 1:100-1:1000* Western blot 1:500-1:3000* Not tested in other applications. *Optimal dilutions/concentrations should be determined by the researcher.Suggested dilutionReferenceIHC (Formalin-fixed paraffin-embedded sections)1:100-1:1000* Western blot1:500-1:3000*	
Comment:	Positive Control: U87-MG	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.77 mg/mL	
Buffer:	0.1M Tris, 0.1M Glycine, 20 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative	
Preservative:	Thimerosal (Merthiolate)	
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw	


cycles.

Images



Immunohistochemistry

Image 1. IHC-P Image Immunohistochemical analysis of paraffin-embedded U374 xenograft, using Frizzled 9, antibody at 1:500 dilution.



Western Blotting

Image 2. WB Image Sample (30 ug of whole cell lysate) A: U87-MG 7.5% SDS PAGE antibody diluted at 1:1000