



[Go to Product page](#)

Datasheet for ABIN2866284
NF-kB p65 ELISA Kit

Overview

Quantity:	96 tests
Target:	NF-kB p65 (NFkBp65)
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA

Product Details

Purpose:	DNA-binding ELISA that facilitate the study of transcription factor activation in mammalian tissue and cell culture extracts.
Brand:	TransAM®
Sample Type:	Cell Extracts, Tissue Samples
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	TransAM NFkB p65 Kits are tested for sensitivity in detecting NFkB activation.
Characteristics:	<p>Transcription factors are DNA-binding proteins that tightly regulate gene expression. They consist of two distinct domains - one that displays high affinity for a specific DNA sequence and one that confers transcriptional activity. Transcription factors are activated by phosphorylation of specific residues or by processing bound inhibitory proteins. Understanding and quantifying transcription factors is essential for the study of cell functions in relation to differentiation, brain activity, immune response, inflammation and various disease states.</p> <p>TransAM® Kits are sensitive, non-radioactive transcription factor ELISA kits that facilitate the</p>

Product Details

study of transcription factor activation in mammalian tissue and cell extracts.

TransAM® Kits are DNA-binding ELISAs that facilitate the study of transcription factor activation in mammalian tissue and cell extracts. Each kit includes a 96-stripwell plate in which multiple copies of a specific double-stranded oligonucleotide have been immobilized. When nuclear or whole-cell extract is added, activated transcription factor of interest binds the oligonucleotide at its consensus binding site and is quantified using the included antibody, which is specific for the bound, active form of the transcription factor being studied.

Components:	One or five 96-well plate(s) with plate sealer(s), primary antibody, HRP-conjugated secondary antibody, wild-type and mutated competitor oligonucleotides, positive control cell extract, DTT, Protease Inhibitor Cocktail, Lysis, Binding, 10X Washing and 10X Antibody Binding Buffers, and Developing and Stop Solutions.
-------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Target Details

Target:	NF-kB p65 (NFkBp65)
Alternative Name:	Nfkb p65 (NFkBp65 Products)
Pathways:	NF-kappaB Signaling , RTK Signaling , TCR Signaling , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Cellular Response to Molecule of Bacterial Origin , Hepatitis C , Toll-Like Receptors Cascades , S100 Proteins

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Nuclear extracts prepared from Jurkat cells that are either unstimulated or stimulated with TPA + CI are diluted to 0.625 µg/well and assayed using the TransAM NFkB p65 Kit. The ratio of the signals from the stimulated over unstimulated cells must be above 4. Lot No. 26914050 was developed for 5 minutes. It gave a ratio of 4.7 (Figure 1). This ratio may vary depending on the basal level of NFkB activation in a given cell type. TransAM NFkB p65 Kits are also tested for specificity in detecting NFkB activation. TransAM NFkB assays are performed in the presence of an excess of oligonucleotide containing a wild-type or mutated NFkB consensus binding site (Figure 2). At 20X excess, the wild-type oligonucleotide prevents NFkB binding to the probe immobilized on the plate. Conversely, the mutated oligonucleotide has no effect on NFkB binding.
Assay Time:	5 h

Application Details

Plate: Pre-coated

Restrictions: For Research Use only

Handling

Storage: 4 °C/-20 °C/-80 °C

Storage Comment: Except for the cell extract that must be kept at -80°C, the kit components can be stored at -20°C prior to first use. Then, we recommend storing the kit at 4°C except for the oligonucleotides, DTT, primary antibody and Protease Inhibitor Cocktail that should be kept at -20°C, and the cell extract at -80°C. This product is guaranteed for 6 months from date of receipt.

Expiry Date: 6 months
