

[Go to Product page](#)

## Datasheet for ABIN2866288

### p53 ELISA Kit

#### Overview

Quantity:	96 tests
Target:	p53 (TP53)
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA

#### Product Details

Purpose:	DNA-binding ELISA that facilitate the study of transcription factor activation in mammalian tissue and cell culture extracts.
Brand:	TransAM®
Sample Type:	Cell Extracts, Tissue Samples
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	TransAM p53 Kits are tested for sensitivity in detecting p53 in nuclear extracts from MCF-7 cells that are either untreated or treated for 3 hours with 200 µM H <sub>2</sub> O <sub>2</sub> .
Characteristics:	Transcription factors are DNA-binding proteins that tightly regulate gene expression. They consist of two distinct domains - one that displays high affinity for a specific DNA sequence and one that confers transcriptional activity. Transcription factors are activated by phosphorylation of specific residues or by processing bound inhibitory proteins. Understanding and quantifying transcription factors is essential for the study of cell functions in relation to differentiation, brain activity, immune response, inflammation and various disease states.

## Product Details

TransAM® Kits are sensitive, non-radioactive transcription factor ELISA kits that facilitate the study of transcription factor activation in mammalian tissue and cell extracts.

TransAM® Kits are DNA-binding ELISAs that facilitate the study of transcription factor activation in mammalian tissue and cell extracts. Each kit includes a 96-stripwell plate in which multiple copies of a specific double-stranded oligonucleotide have been immobilized. When nuclear or whole-cell extract is added, activated transcription factor of interest binds the oligonucleotide at its consensus binding site and is quantified using the included antibody, which is specific for the bound, active form of the transcription factor being studied.

Components:	One or five 96-well plate(s) with plate sealer(s), primary antibody, HRP-conjugated secondary antibody, wild-type and mutated competitor oligonucleotides, positive control cell extract, DTT, Protease Inhibitor Cocktail, Lysis, Binding, 10X Washing and 10X Antibody Binding Buffers, and Developing and Stop Solutions.
-------------	--

## Target Details

Target:	p53 (TP53)
Alternative Name:	p53 ( <a href="#">TP53 Products</a> )
Pathways:	<a href="#">p53 Signaling</a> , <a href="#">MAPK Signaling</a> , <a href="#">PI3K-Akt Signaling</a> , <a href="#">Apoptosis</a> , <a href="#">AMPK Signaling</a> , <a href="#">Chromatin Binding</a> , <a href="#">ER-Nucleus Signaling</a> , <a href="#">Positive Regulation of Endopeptidase Activity</a> , <a href="#">Hepatitis C</a> , <a href="#">Protein targeting to Nucleus</a> , <a href="#">Autophagy</a> , <a href="#">Warburg Effect</a>

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	These extracts are diluted down to 0.625 µg/well and assayed using the TransAM p53 Kit. The ratio of the signals from the treated over the untreated cells must be above 4. Lot 08614019 was developed for 3 minutes. It gave a ratio of 4.3 (Figure 1). The basal level of p53 expression, and this ratio, may vary depending on the cell type tested and the treatment used. TransAM p53 Kits are also tested for specificity in detecting p53 activation. TransAM p53 assays are performed in the presence of an excess of oligonucleotide containing a wild-type or mutated p53 consensus binding site (Figure 2). At 20X excess, the wild-type oligonucleotide prevents p53 binding to the probe immobilized on the plate. Conversely, the mutated oligonucleotide has little effect on p53 binding.
Assay Time:	5 h
Plate:	Pre-coated

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Storage: 4 °C/-20 °C/-80 °C

---

Storage Comment: Except for the cell extract that must be kept at -80°C, the kit components can be stored at -20°C prior to first use. Then, we recommend storing the kit at 4°C except for the oligonucleotides, DTT, poly[d(I-C)], primary antibody and Protease Inhibitor Cocktail that should be kept at -20°C, and the cell extract at -80°C. This product is guaranteed for 6 months from date of receipt.

---

Expiry Date: 6 months