

Datasheet for ABIN2868761

anti-Beclin 1 antibody (C-Term) (Atto 594)[Go to Product page](#)**3** Images

Overview

Quantity:	100 µg
Target:	Beclin 1 (BECN1)
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Beclin 1 antibody is conjugated to Atto 594
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Synthetic peptide from the C-terminus of human Beclin 1
Specificity:	Predicted molecular weight at ~51 kDa.
Cross-Reactivity:	Human, Rat
Purification:	Peptide Affinity Purified

Target Details

Target:	Beclin 1 (BECN1)
Alternative Name:	Beclin 1 (BECN1 Products)
Background:	Beclin 1 is the mammalian ortholog of the yeast autophagy -related gene Atg6. It regulates autophagy and has an important role in development, tumorigenesis and neurodegeneration (1).

Target Details

Beclin 1 is localized within cytoplasmic structures including the mitochondria, yet overexpression reveals some nuclear staining (2). Researchers have found that schizophrénia is associated with low levels of Beclin-1 in the hippocampus (3). It may also protect against infection by a neurovirulent strain of Sindbis virus (4).


Gene ID:	8678
NCBI Accession:	NP_001300927
UniProt:	Q14457
Pathways:	Autophagy

Application Details

Application Notes:	• WB (1:1000) • ICC/IF (1:100) • optimal dilutions for assays should be determined by the user.
Comment:	A 1:1000 dilution of ABIN2868761 was sufficient for detection of Beclin1 on 293T lysates using Goat anti-rabbit IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C




Western Blotting

Image 1. Western blot analysis of Human Embryonic kidney epithelial cell line (HEK293T) lysate showing detection of ~51 kDa Beclin 1 protein using Rabbit Anti-Beclin 1 Polyclonal Antibody (ABIN2868761). Lane 1: MW Ladder. Lane 2: Human 293T (20 µg). Load: 20 µg. Block: 5 % milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-Beclin 1 Polyclonal Antibody (ABIN2868761) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at 1:2000 for 1 hour at RT. Color Development: TMB solution for 12 min at RT. Predicted/Observed Size: ~51 kDa.



Western Blotting

Image 2. Western blot analysis of Human HeLa cell lysates showing detection of ~51kDa Beclin 1 protein using Rabbit Anti-Beclin 1 Polyclonal Antibody . Lane 1: MW Ladder. Lane 2: Human HeLa (20 µg). Load: 20 µg. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-Beclin 1 Polyclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at 1:2000 for 1 hour at RT. Color Development: TMB solution for 12 min at RT. Predicted/Observed Size: ~51kDa.



Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Beclin 1 Polyclonal Antibody . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-Beclin 1 Polyclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Golgi apparatus, Dendrites and Cell bodies of cerebellar Purkinje cells. Magnification: 60X. (A) DAPI (blue) nuclear stain (B)

Phalloidin Texas Red F-Actin stain (C) Beclin 1 Antibody (D)
Composite.