

Datasheet for ABIN2869012

anti-p53 antibody (C-Term) (Atto 594)[Go to Product page](#)**3** Images

Overview

Quantity:	100 µg
Target:	p53 (TP53)
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is conjugated to Atto 594
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Synthetic peptide from the C-terminal of human tumor protein p53.
Specificity:	Predicted molecular weight at ~43.7 kDa. Observed bands at ~53 kDa.
Cross-Reactivity:	Human
Purification:	Peptide Affinity Purified

Target Details

Target:	p53 (TP53)
Alternative Name:	p53 (TP53 Products)
Background:	The p53 protein (tumor protein 53 or TP53) is a DNA-binding cell cycle-regulating transcription factor that governs cell division and the fine balance between cell death and cell survival (1).

Target Details

P53 plays a critical role in tumor suppression and hence it is often described as "the guardian of the genome", "the guardian angel gene", or the "master watchman." This also refers to its role in conserving stability by preventing genome mutation (2). Defects in p53 are linked to >50 % of human cancers, and restoring p53 function to these cancer cells can induce growth arrest and apoptosis (3). When p53 has been damaged, it can also lead to autoimmune disorders (4, 5).

Gene ID: 7157

NCBI Accession: [NP_000537](#)

UniProt: [P04637](#)

Pathways: [p53 Signaling](#), [MAPK Signaling](#), [PI3K-Akt Signaling](#), [Apoptosis](#), [AMPK Signaling](#), [Chromatin Binding](#), [ER-Nucleus Signaling](#), [Positive Regulation of Endopeptidase Activity](#), [Hepatitis C](#), [Protein targeting to Nucleus](#), [Autophagy](#), [Warburg Effect](#)

Application Details

Application Notes:

- WB (1:1000)
- ICC/IF (1:100)
- optimal dilutions for assays should be determined by the user.

Comment: A 1:1000 dilution of ABIN2869012 was sufficient for detection of p53 on HeLa cell lysates using Goat anti-rabbit IgG:HRP as the secondary antibody.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL


Buffer: PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: Conjugated antibodies should be stored at 4°C




Western Blotting

Image 1. Western blot analysis of Human HeLa and HEK293T cell lysates showing detection of ~43.7 kDa p53 protein using Rabbit Anti-p53 Polyclonal Antibody (ABIN2869012). Lane 1: MW Ladder. Lane 2: Human HeLa (20 µg). Lane 3: Human 293T (20 µg). Load: 20 µg. Block: 5 % milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-p53 Polyclonal Antibody (ABIN2869012) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at 1:2000 for 1 hour at RT. Color Development: TMB solution for 12 min at RT. Predicted/Observed Size: ~43.7 kDa.

Immunofluorescence (fixed cells)

Image 2. Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-p53 Polyclonal Antibody . Tissue: Cervical Cancer cell line (HeLa). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-p53 Polyclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: DAPI (blue) nuclear stain at 1:5000 for 5 min RT. Localization: Cytoplasm, PML body, Endoplasmic Reticulum. Magnification: 40X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) p53 Antibody (D) Composite.



Western Blotting

Image 3. Western blot analysis of Human A431 showing detection of ~43.7kDa p53 protein using Rabbit Anti-p53 Polyclonal Antibody . Lane 1: MW Ladder. Lane 2: Human A431 (20 µg). Load: 20 µg. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-p53 Polyclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at 1:2000 for 1 hour at RT. Color Development: TMB solution for 12 min at RT. Predicted/Observed Size: ~43.7kDa.