



[Go to Product page](#)

Datasheet for ABIN2870544

## FCGR3A Protein (AA 17-208) (His tag,AVI tag,Biotin)

### 2 Images

#### Overview

Quantity:	200 µg
Target:	FCGR3A
Protein Characteristics:	AA 17-208
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This FCGR3A protein is labelled with His tag,AVI tag,Biotin.

#### Product Details

Brand:	MABSol@,PrecisionAvi
Sequence:	AA 17-208
Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Characteristics:	This protein carries an Avi tag (Avitag™) at the C-terminus, followed by a polyhistidine tag. The protein has a calculated MW of 25.5 kDa. The protein migrates as 40-55 kDa on a SDS-PAGE gel under reducing (R) condition due to glycosylation.
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

## Target Details

---

Target: FCGR3A

Alternative Name: Fc gamma RIIIA / CD16a ([FCGR3A Products](#))

Background: CD16 is a low affinity Fc receptor, and has been identified as Fc receptors FcγRIIIa (CD16a) and FcγRIIIb (CD16b). These receptors bind to the Fc portion of IgG antibodies. CD16 encoded by two different highly homologous genes in a cell type-specific manner. CD16 is found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16a antigen is also known as Low affinity immunoglobulin gamma Fc region receptor III-A, Fc-gamma RIII-alpha. CD16b is a low-affinity, GPI-linked receptor expressed by neutrophils and eosinophils, whereas CD16a is an intermediate affinity polypeptide-anchored transmembrane glycoprotein expressed natural killer cells, macrophages, subpopulation of T-cells, immature thymocytes and placental trophoblasts. CD16a is involved in phagocytosis, secretion of enzymes and inflammatory mediators, antibody-dependent cytotoxicity and clearance of immune complexes. Aberrant expression or mutations of CD16a is implicated in susceptibility to recurrent viral infections, systemic lupus erythematosus, and alloimmune neonatal neutropenia.

Molecular Weight: 25.5 kDa

## Application Details

---

Comment: Ready-to-use Avitag™ biotinylated protein:  
The product is exclusively produced using the Avitag™ technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA.

This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized


## Handling

Buffer: PBS, pH 7.4

Handling Advice: Please avoid repeated freeze-thaw cycles.


Storage: -20 °C

## Images



### SDS-PAGE

**Image 1.** Biotinylated Human FCGR3A / CD16a (F158) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 92%.



### Binding Studies

**Image 2.** Immobilized Yervoy Ipilimumab at 10µg/mL (100 µl/well), can bind Biotinylated Human FCGR3A / CD16a (F158) (Cat# CDA-H82E8) with a linear of 0.3-10 µg/mL.