

Datasheet for ABIN2870707

**CD137 Protein (AA 24-211) (Fc Tag)****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	CD137 (TNFRSF9)
Protein Characteristics:	AA 24-211
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD137 protein is labelled with Fc Tag.

## Product Details

Sequence:	AA 24-211
Characteristics:	This protein carries a human IgG1 Fc tag at the C-terminus. The protein has a calculated MW of 46.7 kDa. The protein migrates as 50-66 kDa under reducing (R) condition (SDS-PAGE).
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

## Target Details

Target:	CD137 (TNFRSF9)
Alternative Name:	4-1BB ( <a href="#">TNFRSF9 Products</a> )
Background:	4-1BB is also known as CD137, tumor necrosis factor receptor superfamily member 9 (TNFRSF9), induced by lymphocyte activation (ILA), is a co-stimulatory molecule of the tumor necrosis factor (TNF) receptor superfamily. CD137 can be expressed by activated T cells, but to

## Target Details

a larger extent on CD8 than on CD4 T cells. In addition, CD137 expression is found on dendritic cells, follicular dendritic cells, natural killer cells, granulocytes and cells of blood vessel walls at sites of inflammation. The best characterized activity of CD137 is its costimulatory activity for activated T cells. Crosslinking of CD137 enhances T cell proliferation, IL-2 secretion survival and cytolytic activity. Further, it can enhance immune activity to eliminate tumors in mice. CD137 can enhance activation-induced T cell apoptosis when triggered by engagement of the TCR/CD3 complex. In addition, 4-1BB/4-1BBL co-stimulatory pathway has been shown to augment secondary CTL responses to several viruses, and meanwhile augment anti-tumor immunity. 4-1BB thus is a promising candidate for immunotherapy of human cancer. CD137 has been shown to interact with TRAF2.

Molecular Weight:	46.7 kDa
-------------------	----------

NCBI Accession:	<a href="#">NP_001070976</a>
-----------------	------------------------------

Pathways:	<a href="#">Cancer Immune Checkpoints</a>
-----------	---

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------



## Handling

Format:	Lyophilized
---------	-------------

Buffer:	Tris with Glycine, Arginine and NaCl, pH 7.5
---------	--

Handling Advice:	Please avoid repeated freeze-thaw cycles.
------------------	---

Storage:	-20 °C
----------	--------



**ELISA**

**Image 1.** Immobilized Mouse 4-1BB, Fc Tag (ABIN2870706,ABIN2870707) at 2 µg/mL (100 µL/well) can bind Biotinylated Human 4-1BB Ligand, Fc Tag, Avi Tag (ABIN5954977,ABIN6253642) with a linear range of 0.01-0.156 µg/mL (Routinely tested).

**SDS-PAGE**

**Image 2.** Mouse 4-1BB, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 92%.

**ELISA**

**Image 3.** Immobilized Mouse 4-1BB, Fc Tag (ABIN2870706,ABIN2870707) at 2 µg/mL (100 µL/well) can bind Human 4-1BB Ligand, Mouse IgG2a Fc Tag, low endotoxin (ABIN5954978,ABIN6253610) with a linear range of 20-312 ng/mL (QC tested).