antibodies

# Datasheet for ABIN2870711 ADAM17 Protein (AA 18-563) (His tag)





#### Overview

Quantity:	50 µg
Target:	ADAM17
Protein Characteristics:	AA 18-563
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ADAM17 protein is labelled with His tag.

### Product Details

Sequence:	AA 18-563
Characteristics:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of 63.4 kDa. The protein migrates as 19 kDa and 20 kDa(Propeptide),55-66 kDa (Mature-ADAM17) and 75-94 kDa(Pro-ADAM17) under reducing (R) condition (SDS-PAGE) due to glycosylation.
Purity:	>85 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per $\mu$ g by the LAL method.

### Target Details

Target:	ADAM17
Alternative Name:	ADAM17 (ADAM17 Products)
Background:	Disintegrin and metalloproteinase domain-containing protein 17 (ADAM17), a member of the
	ADAM protein family of disintegrins and metalloproteases, is also known as TNF-alpha

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2870711 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	convertase, TNF-alpha-converting enzyme and CD156b, which contains one disintegrin domain
	and one peptidase M12B domain. ADAM17 can cleave the membrane-bound precursor of TNF-
	alpha to its mature soluble form. ADAM17 is also responsible for the proteolytical release of
	soluble JAM3 from endothelial cells surface (By similarity) and proteolytic release of several
	other cell-surface proteins, including p75 TNF-receptor, interleukin 1 receptor type II, p55 TNF-
	receptor, transforming growth factor-alpha, L-selectin, growth hormone receptor, MUC1 and the
	amyloid precursor protein. Furthermore, ADAM17 acts as an activator of Notch pathway by
	mediating cleavage of Notch, generating the membrane-associated intermediate fragment
	called Notch extracellular truncation.
Molecular Weight:	63.4 kDa
UniProt:	Q9Z0F8
Pathways:	Notch Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Response to

## Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	50 mM MES, 150 mM NaCl, pH 6.5
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C

Growth Hormone Stimulus

kDa	М	R
116.0	-	
66.2	-	-
45.0	-	
35.0	-	
25.0	-	
18.4	-	
14.4	-	

#### SDS-PAGE

**Image 1.** Mouse ADAM17, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2870711 | 09/11/2023 | Copyright antibodies-online. All rights reserved.