antibodies.com

# Datasheet for ABIN2870739 FCGR3A Protein (AA 32-215) (His tag)

2 Images



#### Overview

Quantity:	100 µg
Target:	FCGR3A
Protein Characteristics:	AA 32-215
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This FCGR3A protein is labelled with His tag.

### Product Details

Brand:	MABSol®
Sequence:	AA 32-215
Characteristics:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of 22.95 kDa. The protein migrates as 36-45 kDa under reducing (R) condition (SDS-PAGE).
Purity:	>95 % as determined by reduced SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per $\mu$ g by the LAL method.

## Target Details


Target:	FCGR3A
Alternative Name:	Fc gamma RIIIA / CD16a (FCGR3A Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2870739 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

# Target Details

Background:	CD16 is a low affinity Fc receptor, and has been identified as Fc receptors Fc $\gamma$ RIIIa (CD16a) and
	Fc $\gamma$ RIIIb (CD16b). These receptors bind to the Fc portion of IgG antibodies. CD16 encoded by
	two different highly homologous genes in a cell type-specific manner.CD16 is found on the
	surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and
	macrophages. CD16a antigen is also known as Low affinity immunoglobulin gamma Fc region
	receptor III-A, Fc-gamma RIII-alpha. CD16b is a low-affinity, GPI-linked receptor expressed by
	neutrophils and eosinophils, whereas CD16a is an intermediate affinity polypeptide-anchored
	transmembrane glycoprotein expressed natural killer cells, macrophages, subpopulation of T-
	cells, immature thymocytes and placentaltrophoblasts.CD16a is involved in phagocytosis,
	secretion of enzymes and inflammatory mediators, antibody-dependent cytotoxicity and
	clearance of immune complexes. Aberrant expression or mutations of CD16a is implicated in
	susceptibility to recurrent viral infections, systemic lupus erythematosus, and alloimmune
	neonatal neutropenia.
Molecular Weight:	23.0 kDa
Molecular Weight: UniProt:	· · · · · · · · · · · · · · · · · · ·
	23.0 kDa
UniProt:	23.0 kDa
UniProt: Application Details	23.0 kDa Q5D518
UniProt: Application Details	23.0 kDa Q5D518
UniProt: Application Details Restrictions:	23.0 kDa Q5D518
UniProt: Application Details Restrictions: Handling	23.0 kDa Q5D518 For Research Use only
UniProt: Application Details Restrictions: Handling Format:	23.0 kDa Q5D518 For Research Use only Lyophilized

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2870739 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



### 

### SDS-PAGE

**Image 1.** Mouse FCGR3A / CD16a, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

### **Binding Studies**

**Image 2.** Immobilized OKT3 mAb (mouse IgG2a) at 10  $\mu$  g/mL (100  $\mu$ I/well) can bind Mouse FCGR3A / CD16a, His Tag (Cat# CDA-M52H8) with a linear range of 0.16-0.6  $\mu$  g/mL.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2870739 | 09/11/2023 | Copyright antibodies-online. All rights reserved.