antibodies -online.com





IL4 Receptor Protein (AA 26-233) (His tag)

2 Images



Overview

Quantity:	100 μg
Target:	IL4 Receptor (IL4R)
Protein Characteristics:	AA 26-233
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL4 Receptor protein is labelled with His tag.

Product Details

Sequence:	AA 26-233
Characteristics:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of 26.3 kDa. The protein migrates as 37-45 kDa under reducing (R) condition (SDS-PAGE).
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.

Target Details

Target:	IL4 Receptor (IL4R)
Alternative Name:	IL-4 R alpha (IL4R Products)
Background:	IL-4 is a pleiotropic cytokine produced by activated Th2 cells and mast cells, and plays a pivotal role in immune responses. The effects of IL-4 are mediated after binding to high affinity
	receptor complexes present on hematopoietic as well as non-hematopoietic cells.

Hematopoietic cellular responses to IL-4 are mediated by a high affinity receptor complex comprised of the 140 kDa IL4Rα (CD124)subunit and the 70 kDa common cytokine γc chain (CD132). Interleukin 4 Receptor (IL4R) also known as CD124, IL4Rα and BSF receptor, is a type I cytokine receptor produced by activated Th2 cells and mast cells, and plays an important role in Th2--biased immune responses, alternative macrophage activation, mucosal immunity, allergic inflammation, tumor progression, and atherogenesis. A soluble form of the encoded IL4R protein can be produced by an alternate splice variant or by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. IL4R can alternatively associate with IL-

13Ra1 to form the type II receptor which is responsive to both IL-4 and IL-13. Interleukin-4 receptor has been shown to interact with SHC1.

Molecular Weight: 26.3 kDa

NCBI Accession: NP_001008700


Pathways: JAK-STAT Signaling, Positive Regulation of Immune Effector Process


Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C





ELISA

Image 1. Immobilized IL-4 Protein, Mouse, Recombinant at $2 \mu g/mL$ (100 $\mu L/well$) can bind Mouse IL-4 R alpha, His Tag (ABIN2870747,ABIN2870748) with a linear range of 0.039-1.25 $\mu g/mL$ (Routinely tested).

SDS-PAGE

Image 2. Mouse IL-4 R alpha, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.