

Datasheet for ABIN2872465

**anti-Urotensin 2 antibody (Preproprotein)**[Go to Product page](#)

## Overview

|                      |                                            |
|----------------------|--------------------------------------------|
| Quantity:            | 100 µL                                     |
| Target:              | Urotensin 2 (UTS2)                         |
| Binding Specificity: | AA 41-68, Preproprotein                    |
| Reactivity:          | Human                                      |
| Host:                | Rabbit                                     |
| Clonality:           | Polyclonal                                 |
| Conjugate:           | This Urotensin 2 antibody is un-conjugated |
| Application:         | Immunohistochemistry (IHC)                 |

## Product Details

|               |                                                  |
|---------------|--------------------------------------------------|
| Immunogen:    | 5-10 mg of > 70% pure peptide conjugated to KLH. |
| Purification: | unpurified serum                                 |

## Target Details

|                   |                                                          |
|-------------------|----------------------------------------------------------|
| Target:           | Urotensin 2 (UTS2)                                       |
| Alternative Name: | Urotensin II ( <a href="#">UTS2 Products</a> )           |
| Pathways:         | <a href="#">Negative Regulation of Hormone Secretion</a> |

## Application Details

|                    |                                                                    |
|--------------------|--------------------------------------------------------------------|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--------------------------------------------------------------------|

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Storage: 4 °C,-20 °C

Storage Comment: Store antibodies at 2-4°C before and at -20°C after rehydration. The product is stable for approximately 3-6 weeks after rehydration at -20°C or below.