

Datasheet for ABIN2879101
anti-c-MET antibody (AA 1316-1365)



[Go to Product page](#)

2 Images

Overview

Quantity:	50 µL
Target:	c-MET (MET)
Binding Specificity:	AA 1316-1365
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This c-MET antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic peptide from human Met (aa1316-1365). Type of Immunogen: Synthetic peptide
Isotype:	IgG
Specificity:	Met (Ab-1349) Antibody detects endogenous levels of total Met protein.
Purification:	Immunoaffinity purified

Target Details

Target:	c-MET (MET)
---------	-------------

Target Details


Alternative Name:	c-Met (MET Products)
Background:	Name/Gene ID: MET Subfamily: HGF Receptor/MET Family: Protein Kinase Synonyms: MET, AUTS9, C-Met, HGF/SF receptor, HGFR, Met kinase, Proto-oncogene c-Met, Scatter factor receptor, SF receptor, HGF receptor, RCCP2, Tyrosine-protein kinase Met
Gene ID:	4233
UniProt:	P08581
Pathways:	RTK Signaling , Carbohydrate Homeostasis , Synaptic Membrane , Signaling of Hepatocyte Growth Factor Receptor

Application Details

Application Notes:	Approved: IHC, IHC-P (1:100), Peptide-ELISA (1:10000), WB (1:500 - 1:1000)
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS (without Mg ²⁺ , Ca ²⁺), pH 7.4, 150 mM sodium chloride, 0.02 % sodium azide, 50 % glycerol
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	avoid freeze thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze-thaw cycles.



Immunohistochemistry

Image 1. Anti-c-Met antibody IHC staining of human small intestine. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1100321 dilution 10 ug/ml.



Immunohistochemistry

Image 2. Anti-c-Met antibody IHC staining of human liver. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1100321 dilution 10 ug/ml.