

Datasheet for ABIN288463 anti-Resistin antibody (Biotin)



_					
	W	0	rv	10	W

Overview			
Quantity:	50 μg		
Target:	Resistin (RETN)		
Reactivity:	Human		
Host:	Goat		
Clonality:	Polyclonal		
Conjugate:	This Resistin antibody is conjugated to Biotin		
Application:	Western Blotting (WB), ELISA		
Product Details			
Immunogen:	Resistin antibody (biotin) was raised in goat using highly pure recombinant human resistinas		
	the immunogen.		
Target Details			
Target:	Resistin (RETN)		
Alternative Name:	Resistin (RETN Products)		
Background:	Resistin belongs to a family of tissue-specific cytokines termed FIZZ (found in inflammatory		
	zones) and RELM. The three known members of this family, Resistin, RELM-alpha and RELM-		
	beta share a highly conserved C-terminal domain, characterized by 10 cysteine residues with a		
	unique spacing motif of C-X11-C-X8-C-X-C-X3-C-X10-C-X-C-X9-C-C. Resistin is an adipose-		
	derived cytokine (adipokine) whose physiological function and molecular targets are largely		
	unknown. Studies have shown that Resistin suppresses insulin's ability to stimulate glucose		
	The state of the s		

uptake, and postulated that Resistin might be an important link between obesity and Type 2

Target Details

l arget Details		
	diabetes. Other studies have indicated that Resistin expression is severely suppressed in	
	obesity and that it may act as a feedback regulator of Adipogenesis.	
Molecular Weight:	20.2 kDa (predicted detection band MW)	
Pathways:	Feeding Behaviour, Smooth Muscle Cell Migration	
Application Details		
Application Notes:	ELISA: 0.25-1 μg/mL, WB: 0.1-0.2 μg/mL	
	Optimal conditions should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Reconstitute in sterile PBS with 0.1 % BSA to a concentration of at least 50 µg/mL	
Concentration:	Lot specific	
Buffer:	Lyophilized from PBS, pH 7.2.	
Handling Advice:	Avoid repeated freeze/thaw cycles.	
	Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at -20 °C until reconstitution. Following reconstitution product may be stored at 4 °C in	

the short term. For long term storage aliquot and freeze at -20 $^{\circ}\text{C}.$