

Datasheet for ABIN2891611

anti-TGFB1 antibody (Tyr369)



Overview

Overview	
Quantity:	100 μL
Target:	TGFB1
Binding Specificity:	Tyr369
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TGFB1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunofluorescence (IF)
Product Details	
Immunogen:	Synthetic peptide around Tyr369 of human TGFB1 / TGF Beta
	Type of Immunogen: Synthetic peptide
Specificity:	Human TGFB1 / TGF Beta
Purification:	Immunoaffinity purified
Target Details	
Target:	TGFB1
Alternative Name:	TGFB1 / TGF Beta 1 (TGFB1 Products)
Background:	Name/Gene ID: TGFB1

Target Details

	Family: TGF beta
	Synonyms: TGFB1, CED, DPD1, Diaphyseal dysplasia 1, LAP, Latency-associated peptide, TGF Beta, TGFbeta, TGF-beta 1 protein, TGF-beta-1, TGFB
Gene ID:	7040
UniProt:	P01137
Pathways:	EGFR Signaling Pathway, Dopaminergic Neurogenesis, Cellular Response to Molecule of Bacterial Origin, Glycosaminoglycan Metabolic Process, Regulation of Leukocyte Mediated Immunity, Regulation of Muscle Cell Differentiation, Positive Regulation of Immune Effector Process, Cell-Cell Junction Organization, Production of Molecular Mediator of Immune Response, Ribonucleoside Biosynthetic Process, Skeletal Muscle Fiber Development, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy, Cancer Immune Checkpoints

Application Details

Application Notes:

Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.2, 0.05 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Optimal working dilution should be determined by the investigator.