

Datasheet for ABIN289532

anti-Resistin antibody



_					
	W	0	rv	10	W

Quantity:	100 μg		
Target:	Resistin (RETN)		
Reactivity:	Mouse		
Host:	Goat		
Clonality:	Polyclonal		
Conjugate:	This Resistin antibody is un-conjugated		
Application:	Western Blotting (WB), ELISA, Blocking Antibody (Inhibition)		
Product Details			
Immunogen:	Resistin antibody was raised in goat using highly pure recombinant murine resistin as the		
	immunogen.		
Purification:	Purified		
Target Details			
Target:	Resistin (RETN)		
Alternative Name:	Resistin (RETN Products)		
Background:	Resistin belongs to a family of tissue-specific cytokines termed FIZZ (found in inflammatory		
	zones) and RELM. The three known members of this family, Resistin, RELM-alpha and RELM-		
	beta share a highly conserved C-terminal domain, characterized by 10 cysteine residues with a		
	unique spacing motif of C-X11-C-X8-C-X-C-X3-C-X10-C-X-C-X9-C-C. Resistin is an adipose-		
	derived cytokine (adipokine) whose physiological function and molecular targets are largely		

Target Details

l arget Details		
	unknown. Studies have shown that Resistin suppresses insulin's ability to stimulate glucose uptake, and postulated that Resistin might be an important link between obesity and Type 2 diabetes. Other studies have indicated that Resistin expression is severely suppressed in obesity and that it may act as a feedback regulator of Adipogenesis.	
Molecular Weight:	20.2 kDa (predicted detection band MW)	
Pathways:	Feeding Behaviour, Smooth Muscle Cell Migration	
Application Details		
Application Notes:	Optimal conditions should be determined byt he investigator.	
Restrictions:	For Research Use only	

Handling

Format:	Lyophilized	
Reconstitution:	Reconstitute in sterile PBS with 0.1 % BSA to a concentration of at least 50 µg/mL.	
Concentration:	Lot specific	
Buffer:	Lyophilized from PBS, pH 7.2.	
Handling Advice:	Avoid repeated freeze/thaw cycles.	
	Dilute only prior to immediate use.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at -20 °C until reconstitution. Following reconstitution product may be stored at 4 °C in	
	the short term. For long term storage aliquot and freeze at -20 °C.	