antibodies - online.com







STAT6 Protein






| <i>ا</i> ۱ | 1 | - | M | /1 | 0 | |
|------------|-----|----------|-----|-----|--------|------|
| u | 1// | \vdash | 1 \ | / I | - | \/\/ |
| \sim | ٧. | \sim | r٧ | | \sim | v v |

| Quantity: | 100 μg | | |
|-------------------|--|--|--|
| Target: | STAT6 | | |
| Origin: | Rat | | |
| Source: | Escherichia coli (E. coli) | | |
| Protein Type: | Recombinant | | |
| Application: | Western Blotting (WB), SDS-PAGE (SDS), ELISA, Immunoprecipitation (IP) | | |
| Product Details | | | |
| Characteristics: | Rat Signal Transducer And Activator Of Transcription 6 (STAT6) Protein | | |
| Purity: | > 95 % | | |
| Target Details | | | |
| Target: | STAT6 | | |
| Alternative Name: | Signal Transducer And Activator Of Transcription 6 (STAT6 Products) | | |
| Molecular Weight: | 35.1 kDa | | |
| UniProt: | Q1KQ07 | | |
| Pathways: | JAK-STAT Signaling, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response | | |

Application Details

| Comment: | Please note that the majority of this suppliers proteins are partial length rather than full length. We recommend customers to inquire. |
|--------------------|---|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 200 μg/mL |
| Buffer: | Prior to lyophilization: PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin-300. |
| Preservative: | Dithiothreitol (DTT), ProClin |
| Precaution of Use: | This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only. |
| Storage: | 4 °C,-80 °C |
| Storage Comment: | Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles. |

Images



SDS-PAGE

Image 1. SDS-PAGE analysis of Rat STAT6 Protein.