
antibodies -online.com







anti-Histone 3 antibody (H3K36me3)

Images



Publications



()	11	\sim	rv		۱ ۸
	1 \ /	┙	I \/	╙	1/1

Quantity:	100 μL	
Target:	Histone 3 (H3)	
Binding Specificity:	H3K36me3	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Histone 3 antibody is un-conjugated	
Application:	ication: Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Chromat Immunoprecipitation (ChIP), Immunoprecipitation (IP), ChIP DNA-Sequencing (ChIP-seq)	
Product Details		

Immunogen:	A synthetic methylated peptide corresponding to residues surrounding K36 of human histone H3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Methylated Antibodies
Purification:	Affinity purification

Target Details

Target: Histone 3 (H3)


Target Details

Alternative Name:	Histone H3 (H3 Products)		
Background:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the		
	chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA		
	wrapped around a histone octamer composed of pairs of each of the four core histones (H2A,		
	H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker		
	histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures.		
	This gene is intronless and encodes a replication-dependent histone that is a member of the		
	histone H3 family. Transcripts from this gene lack polyA tails, instead, they contain a		
	palindromic termination element. This gene is located separately from the other H3 genes that		
	are in the histone gene cluster on chromosome 6p22-		
	p21.3.,H3.4,H3/g,H3FT,H3t,HIST3H3,Histone H3,HIST1H3A,Signal Transduction,MAPK-Erk		
	Signaling Pathway,MAPK-P38 Signaling Pathway,Epigenetics & Nuclear Signaling,Epigenetic		
	Modifications,Methylation,Histone H3		
Molecular Weight:	15 kDa		
Gene ID:	8290		
UniProt:	Q16695		
Application Details			
Application Notes:	WB,1:500 - 1:2000,IHC,1:50 - 1:200,IF,1:50 - 1:200,IP,1:50 - 1:200,ChIP,1:20 - 1:100,ChIP-seq,1:20		
	- 1:100		
Restrictions:	For Research Use only		
Handling			
Buffer:	PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
0.	-20 °C		
Storage:			

Product cited in:


Deci, Ferguson, Scatigno, Nguyen: "Modulating Macrophage Polarization through CCR2 Inhibition and Multivalent Engagement." in: **Molecular pharmaceutics**, Vol. 15, Issue 7, pp. 2721-2731, (2018) (PubMed).

Images




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded Human breast cancer using TriMethyl-Histone H3-K36 Rabbit pAb (ABIN3016038, ABIN3016039, ABIN3016040, ABIN1680217 and ABIN6219525) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry

Image 2.



Western Blotting

Image 3.

J	
	Please check the product details page for more images. Overall 15 images are available for ABIN3016039.