

Datasheet for ABIN3017896  
**anti-RAP1GAP antibody**



[Go to Product page](#)

2 Images

## Overview

Quantity:	500 µL
Target:	RAP1GAP
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This RAP1GAP antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

## Product Details

Immunogen:	A synthesized peptide derived from human Rap1GAP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

## Target Details

Target:	RAP1GAP
Alternative Name:	RAP1GAP ( <a href="#">RAP1GAP Products</a> )
Background:	This gene encodes a type of GTPase-activating-protein (GAP) that down-regulates the activity of the ras-related RAP1 protein. RAP1 acts as a molecular switch by cycling between an

## Target Details

inactive GDP-bound form and an active GTP-bound form. The product of this gene, RAP1GAP, promotes the hydrolysis of bound GTP and hence returns RAP1 to the inactive state whereas other proteins, guanine nucleotide exchange factors (GEFs), act as RAP1 activators by facilitating the conversion of RAP1 from the GDP- to the GTP-bound form. In general, ras subfamily proteins, such as RAP1, play key roles in receptor-linked signaling pathways that control cell growth and differentiation. RAP1 plays a role in diverse processes such as cell proliferation, adhesion, differentiation, and embryogenesis. Alternative splicing results in multiple transcript variants encoding distinct proteins. [provided by RefSeq, Aug 2011],RAP1GA1, RAP1GAP1, RAP1GAPII, RAPGAP,Cancer,Epigenetics & Nuclear Signaling,Signal Transduction,Tumor suppressors,RAP1GAP

Molecular Weight: 95 kDa

Gene ID: 5909

UniProt: [P47736](#)

## Application Details

Application Notes: WB,1:500 - 1:2000,IF,1:50 - 1:200

Restrictions: For Research Use only

## Handling


Buffer: PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.



Immunofluorescence

**Image 1.** Immunofluorescence analysis of NIH-3T3 cells using Rap1GAP Rabbit mAb (ABIN1680949, ABIN3017895, ABIN3017896 and ABIN7101534) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Western Blotting

**Image 2.** Western blot analysis of extracts of various cell lines, using Rap1GAP Rabbit mAb (ABIN1680949, ABIN3017895, ABIN3017896 and ABIN7101534) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.