

Datasheet for ABIN302018
anti-CD11c antibody (FITC)[Go to Product page](#)

4 Images

5 Publications

Overview

| | |
|--------------|---|
| Quantity: | 100 tests |
| Target: | CD11c (ITGAX) |
| Reactivity: | Human, Monkey |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This CD11c antibody is conjugated to FITC |
| Application: | Flow Cytometry (FACS) |

Product Details

| | |
|-----------------------------|---|
| Immunogen: | Dendritic cells of synovial fluid |
| Clone: | BU15 |
| Isotype: | IgG1 |
| Specificity: | The antibody BU15 reacts with an extracellular epitope of CD11c (alphaX, p150), a 150 kDa integrin expressed mainly on dendritic cells and tissue macrophages. |
| Cross-Reactivity (Details): | Human, Monkey |
| Purification: | Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |

Target Details

| | |
|---------|---------------|
| Target: | CD11c (ITGAX) |
|---------|---------------|

Target Details

| | |
|-------------------|--|
| Alternative Name: | CD11c (ITGAX Products) |
| Background: | Integrin subunit alpha X,CD11c (p150, alphaX integrin subunit) forms complex with CD18 (beta2 integrin subunit) and is expressed mainly on tissue macrophages and dendritic cells. CD11c binds to complement fragment iC3b, fibrinogen, VCAM-1 and ICAM-2 or e.g. CD90. Like other beta2 integrins, CD11c/CD18 plays roles in cell migration and phagocytosis. Moreover, interaction of CD11c/CD18 with plasminogen regulates plasmin activities, and interaction with heparin counteracts binding of iC3b.,Integrin alpha X, ITGAX, SLEB6 |
| Gene ID: | 3687 |
| UniProt: | P20702 |
| Pathways: | Complement System , Activated T Cell Proliferation , Integrin Complex |

Application Details


| | |
|--------------------|---|
| Application Notes: | Flow cytometry: The reagent is designed for analysis of human blood cells using 20 µL reagent / 100 µL of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests. |
| Comment: | The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary. |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Reconstitution: | No reconstitution is necessary. |
| Buffer: | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Do not freeze. Avoid prolonged exposure to light. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. |


- Product cited in: Angel, Lala, Chen, Edgar, Ostrovsky, Dunbar: "CD14+ antigen-presenting cells in human dermis are less mature than their CD1a+ counterparts." in: **International immunology**, Vol. 19, Issue 11, pp. 1271-9, (2007) ([PubMed](#)).
- Sadhu, Hendrickson, Dick, Potter, Staunton: "Novel tools for functional analysis of CD11c: activation-specific, activation-independent, and activating antibodies." in: **Journal of immunoassay & immunochemistry**, Vol. 29, Issue 1, pp. 42-57, (2007) ([PubMed](#)).
- Myou, Zhu, Boetticher, Qin, Myo, Meliton, Lambertino, Munoz, Hamann, Leff: "Regulation of adhesion of AML14.3D10 cells by surface clustering of beta2-integrin caused by ERK-independent activation of cPLA2." in: **Immunology**, Vol. 107, Issue 1, pp. 77-85, (2002) ([PubMed](#)).
- Van der Vieren, Le Trong, Wood, Moore, St John, Staunton, Gallatin: "A novel leukointegrin, alpha d beta 2, binds preferentially to ICAM-3." in: **Immunity**, Vol. 3, Issue 6, pp. 683-90, (1996) ([PubMed](#)).
- Hogg, Takacs, Palmer, Selvendran, Allen: "The p150,95 molecule is a marker of human mononuclear phagocytes: comparison with expression of class II molecules." in: **European journal of immunology**, Vol. 16, Issue 3, pp. 240-8, (1986) ([PubMed](#)).

Images



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD11c (BU15) FITC antibody (20 µL reagent / 100 µL of peripheral whole blood).



Flow Cytometry

Image 2. Surface staining of human peripheral blood cells with anti-CD11c (BU15) PerCP (monocyte gate).

Flow Cytometry

Image 3. Surface staining of human peripheral blood cells with anti-CD11c (BU15) FITC.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN302018.