antibodies -online.com







anti-CRH antibody (AA 25-194)





| \sim | | | | | |
|--------|-----|----|-----|----|---|
| | 1// | ⊃r | V/I | Φ\ | Λ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | CRH | |
| Binding Specificity: | AA 25-194 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CRH antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC) | |
| Product Details | | |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 25-194 of | |
| | human CRH (NP_000747.1). | |
| Sequence: | LLSRGPVPGA RQAPQHPQPL DFFQPPPQSE QPQQPQARPV LLRMGEEYFL RLGNLNKSPA | |
| | APLSPASSLL AGGSGSRPSP EQATANFFRV LLQQLLLPRR SLDSPAALAE RGARNALGGH | |
| | QEAPERERRS EEPPISLDLT FHLLREVLEM ARAEQLAQQA HSNRKLMEII | |
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Characteristics: | Polyclonal Antibodies | |
| Purification: | Affinity purification | |

Target Details

| l arget Details | Target Details | | |
|---------------------|--|--|--|
| Target: | CRH | | |
| Alternative Name: | CRH (CRH Products) | | |
| Background: | This gene encodes a member of the corticotropin-releasing factor family. The encoded | | |
| | preproprotein is proteolytically processed to generate the mature neuropeptide hormone. In | | |
| | response to stress, this hormone is secreted by the paraventricular nucleus (PVN) of the | | |
| | hypothalamus, binds to corticotropin releasing hormone receptors and stimulates the release | | |
| | of adrenocorticotropic hormone from the pituitary gland. Marked reduction in this protein has | | |
| | been observed in association with Alzheimer's disease. Autosomal recessive hypothalamic | | |
| | corticotropin deficiency has multiple and potentially fatal metabolic consequences including | | |
| | hypoglycemia and hepatitis. In addition to production in the hypothalamus, this protein is also | | |
| | synthesized in peripheral tissues, such as T lymphocytes, and is highly expressed in the | | |
| | placenta. In the placenta it is a marker that determines the length of gestation and the timing of | | |
| | parturition and delivery. A rapid increase in circulating levels of the hormone occurs at the onse | | |
| | of parturition, suggesting that, in addition to its metabolic functions, this protein may act as a | | |
| | trigger for parturition.,CRH,CRF,CRH1,Signal Transduction,Cell Biology & Developmental | | |
| | Biology,Growth factor,Endocrine & Metabolism,Neuroscience,CRH | | |
| Molecular Weight: | 21 kDa | | |
| Gene ID: | 1392 | | |
| UniProt: | P06850 | | |
| Pathways: | Positive Regulation of Peptide Hormone Secretion, Hormone Activity, Negative Regulation of | | |
| | Hormone Secretion, cAMP Metabolic Process, Myometrial Relaxation and Contraction, Feeding | | |
| | Behaviour | | |
| Application Details | | | |
| Application Notes: | WB,1:500 - 1:2000,IHC,1:50 - 1:200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Buffer: | PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3. | | |
| | | | |


Sodium azide


Preservative:

Handling

| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
|--------------------|--|--|
| Handling Advice: | Avoid freeze / thaw cycles | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Avoid freeze / thaw cycles. | |

Images





Western Blotting

Image 1. Western blot analysis of extracts of various cell lines, using CRH antibody (ABIN3021479, ABIN3021480, ABIN3021481 and ABIN6215115) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 μg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.

Western Blotting

Image 2.