antibodies .- online.com





anti-TRADD antibody





Go to Product page

\sim							
	1//	\Box	r۱	/ [\bigcirc	1	٨,

Quantity:	100 μL
Target:	TRADD
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRADD antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant protein of mouse TRADD
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Characteristics:	Polyclonal Antibodies

Target Details

Target:	TRADD
Alternative Name:	TRADD (TRADD Products)
Background:	The protein encoded by this gene is a death domain containing adaptor molecule that interacts with TNFRSF1A/TNFR1 and mediates programmed cell death signaling and NF-kappaB activation. This protein binds adaptor protein TRAF2, reduces the recruitment of inhibitor-of-apoptosis proteins (IAPs) by TRAF2, and thus suppresses TRAF2 mediated apoptosis. This

	protein can also interact with receptor TNFRSF6/FAS and adaptor protein FADD/MORT1, and is involved in the Fas-induced cell death pathway.,TRADD,Hs.89862,Cancer,Invasion and Metastasis,Signal Transduction,MAPK-JNK Signaling Pathway,MAPK-P38 Signaling Pathway,Cell Biology & Developmental Biology,Apoptosis,Death receptors & ligands,Inhibition of Apoptosis,Growth factor,TNF,Death Receptor Signaling Pathway,Immunology & Inflammation,NF-kB Signaling Pathway,Cardiovascular,TRADD
Molecular Weight:	34 kDa
Gene ID:	71609
UniProt:	Q15628

Application Details

Application Notes:	WB,1:500 - 1:2000,IHC,1:50 - 1:100,IF,1:50 - 1:100
Restrictions:	For Research Use only


Endopeptidase Activity, Hepatitis C


NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, Positive Regulation of


Handling

Pathways:

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid freeze / thaw cycles
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.







Western Blotting

Image 1. Western blot analysis of extracts of various cell lines, using TRADD antibody (ABIN3021541, ABIN3021542, ABIN3021543 and ABIN6215219) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using TRADD antibody.

Immunohistochemistry

Image 3.