

Datasheet for ABIN3025667 **anti-HLA-DPB1 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HLA-DPB1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HLA-DPB1 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Non-T, non-B human acute lymphoblastic leukemia REH6 cell line was used as the immunogen for the HLA-DPB1 antibody.
Clone:	BraFB6
Isotype:	IgG2b kappa
Purification:	Protein G affinity chromatography

Target Details

Target:	HLA-DPB1
Alternative Name:	HLA-DPB1 (MHC II) (HLA-DPB1 Products)
Background:	Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35 kDa) and beta (27-30 kDa)

Target Details

chains. It reportedly reacts with B- & non-T, non-B cell lines but not with T- and myeloid cell lines and leukemias. Differential expression of MHC class II antigens on fetal and adult lymphocytes, malignant B cells appears to reflect the stage of cell differentiation which may be useful in the study of lymphoproliferative disorders.

Pathways:	TCR Signaling , Cancer Immune Checkpoints , Human Leukocyte Antigen (HLA) in Adaptive Immune Response
-----------	---

Application Details

Application Notes:	Optimal dilution of the HLA-DPB1 antibody should be determined by the researcher. 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10 mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.\. Flow Cytometry: 0.5-1 µg/million cells in 0.1ml,Immunofluorescence: 0.5-1 µg/mL,Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 min at RT (1)
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Concentration:	0.2 mg/mL
Buffer:	0.2 mg/mL in 1X PBS with 0.1 mg/mL BSA (US sourced) and 0.05 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store the HLA-DPB1 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).