

Datasheet for ABIN3025768
anti-CD11c antibody (AA 600-850)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	CD11c (ITGAX)
Binding Specificity:	AA 600-850
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD11c antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	A recombinant fragment (191 amino acid residues between aa 600-850) from the human protein was used as the immunogen for the CD11c antibody.
Clone:	ITGAX-1242
Isotype:	IgG kappa
Purification:	Protein G affinity chromatography

Target Details

Target:	CD11c (ITGAX)
Alternative Name:	CD11c (ITGAX Products)

Target Details


Background:	Recognizes a protein of 145 kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (MAC-1) and CD11d (ITGAD), but has a unique alpha chain. CD11c has been shown to play a role in phagocytosis, cell migration, and cytokine production by monocytes/macrophages as well as induction of T-cell proliferation by Langerhans cells. CD11c is expressed prominently on the plasma membranes of monocytes, tissue macrophages, NK cells, and most dendritic cells (DCs). A lower level of expression is also observed on neutrophils as a result of its high level of expression on most DCs. An antibody to CD11c may aid in identification of lesions with histiocytic origin. It may also been used as a marker for hairy cell leukemia in paraffin-embedded tissues.
Pathways:	Complement System , Activated T Cell Proliferation , Integrin Complex

Application Details

Application Notes:	Optimal dilution of the CD11c antibody should be determined by the researcher. 1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10 mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.\. Flow Cytometry: 0.5-1 µg/million cells in 0.1ml,Immunofluorescence: 0.5-1 µg/mL,Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 min at RT (1)
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	1 mg/mL in 1X PBS, BSA free, sodium azide free
Preservative:	Azide free
Storage:	4 °C,-20 °C
Storage Comment:	Store the CD11c antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).



Immunohistochemistry

Image 1. IHC testing of human tonsil with CD11c antibody (clone ITGAX/1242). Note specific staining of dendritic cells. Required HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH 9, for 10-20 min.