



## Datasheet for ABIN3026683 anti-Pan Muscle Actin antibody



[Go to Product page](#)

### 6 Images

#### Overview

Quantity:	100 µg
Target:	Pan Muscle Actin
Reactivity:	Human, Rabbit, Rat
Host:	Mouse
Clonality:	Monoclonal
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunofluorescence (IF), Western Blotting (WB)

#### Product Details

Immunogen:	An amino acid sequence common among human muscle actins was used as the immunogen for the pan Muscle Actin antibody.
Clone:	MSA-953
Isotype:	IgG1 kappa
Characteristics:	This antibody recognizes actin of skeletal, cardiac, and smooth muscle cells. It is not reactive with other mesenchymal cells except for myoepithelium. Actin can be resolved on the basis of its isoelectric points into three distinctive components: alpha, beta, and gamma in order of increasing isoelectric point. Anti-muscle specific actin recognizes alpha and gamma isotypes of all muscle groups. Non-muscle cells such as vascular endothelial cells and connective tissues are non-reactive. Also, neoplastic cells of non-muscle-derived tissue such as carcinomas, melanomas, and lymphomas are negative. It stains tumors of smooth muscle (leiomyomas and leiomyosarcomas) as well as skeletal muscle (rhabdomyomas and rhabdomyosarcomas).
Purification:	Protein G affinity chromatography

## Target Details

---

Target:	Pan Muscle Actin
Background:	This antibody recognizes actin of skeletal, cardiac, and smooth muscle cells. It is not reactive with other mesenchymal cells except for myoepithelium. Actin can be resolved on the basis of its isoelectric points into three distinctive components: alpha, beta, and gamma in order of increasing isoelectric point. Anti-muscle specific actin recognizes alpha and gamma isotypes of all muscle groups. Non-muscle cells such as vascular endothelial cells and connective tissues are non-reactive. Also, neoplastic cells of non-muscle-derived tissue such as carcinomas, melanomas, and lymphomas are negative. It stains tumors of smooth muscle (leiomyomas and leiomyosarcomas) as well as skeletal muscle (rhabdomyomas and rhabdomyosarcomas).

## Application Details

---


Application Notes:	Optimal dilution of the pan Muscle Actin antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.\. Flow Cytometry: 0.5-1 µg/million cells in 0.1ml,Immunofluorescence: 0.5-1 µg/mL,Western blot: 0.5-1 µg/mL,Immunohistochemistry (FFPE): 0.5-1 µg/mL for 30 min at RT (1),Prediluted format: incubate for 30 min at RT (2)
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling


---

Concentration:	1 mg/mL
Buffer:	1 mg/mL in 1X PBS, BSA free, sodium azide free
Preservative:	Azide free
Storage:	4 °C,-20 °C
Storage Comment:	Store the pan Muscle Actin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).



**Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)**

**Image 1.** Formalin-fixed, paraffin-embedded human rhabdomyosarcoma stained with pan Muscle Actin antibody (MSA/953)



**Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)**

**Image 2.** Formalin-fixed, paraffin-embedded rat lung stained with pan Muscle Actin antibody



**Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)**

**Image 3.** Formalin-fixed, paraffin-embedded rat skeletal muscle stained with pan Muscle Actin antibody

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN3026683.