

Datasheet for ABIN3029317
anti-Tuberin antibody (pSer1096)



[Go to Product page](#)

2 Images

Overview

Quantity:	0.4 mL
Target:	Tuberin (TSC2)
Binding Specificity:	pSer1096
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tuberin antibody is un-conjugated
Application:	ELISA, Dot Blot (DB)

Product Details

Immunogen:	This phospho-TSC2 antibody was produced from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding pS1096 of human TSC2.
Isotype:	Ig Fraction
Purification:	Antigen affinity purified

Target Details

Target:	Tuberin (TSC2)
Alternative Name:	TSC2 (TSC2 Products)
Background:	Mutations in this gene lead to tuberous sclerosis complex. Its gene product is believed to be a tumor suppressor and is able to stimulate specific GTPases. The protein associates with

Target Details


	hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin. Alternative splicing results in multiple transcript variants encoding different isoforms.
UniProt:	P49815
Pathways:	RTK Signaling , AMPK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Regulation of Cell Size , Tube Formation , Protein targeting to Nucleus

Application Details

Application Notes:	Titration of the phospho-TSC2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Dot blot: 1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	In 1X PBS, pH 7.4, with 0.09 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Aliquot the phospho-TSC2 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.



Dot Blot

Image 1. Dot blot analysis of phospho-TSC2 antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.



Dot Blot

Image 2. Dot blot analysis of phospho-TSC2 antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.