

Datasheet for ABIN3029322
anti-Tuberin antibody (pSer1387)



[Go to Product page](#)

3 Images

Overview

Quantity:	0.4 mL
Target:	Tuberin (TSC2)
Binding Specificity:	pSer1387
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tuberin antibody is un-conjugated
Application:	ELISA, Dot Blot (DB), Immunofluorescence (IF)

Product Details

Immunogen:	This phospho-TSC2 antibody was produced from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding pS1387 of human TSC2.
Isotype:	Ig Fraction
Cross-Reactivity (Details):	Expected species reactivity: Mouse, Rat
Purification:	Antigen affinity purified

Target Details

Target:	Tuberin (TSC2)
Alternative Name:	TSC2 (TSC2 Products)

Target Details

Background: Mutations in TSC2 lead to tuberous sclerosis complex. The protein is believed to be a tumor suppressor and is able to specifically stimulate the intrinsic GTPase activity of the Ras-related protein RAP1A and RAB5. The protein associates with hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin. TSC2 may have a function in vesicular transport, but may also play a role in the regulation of cell growth arrest and in the regulation of transcription mediated by steroid receptors. Interaction between TSC1 and TSC2 may facilitate vesicular docking.

UniProt: [P49815](#)

Pathways: [RTK Signaling](#), [AMPK Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Regulation of Cell Size](#), [Tube Formation](#), [Protein targeting to Nucleus](#)

Application Details

Application Notes: Titration of the phospho-TSC2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.\. Dot blot: 1:500,Immunofluorescence: 1:10-1:50

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: In 1X PBS, pH 7.4, with 0.09 % sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Aliquot the phospho-TSC2 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.




Immunofluorescence

Image 1. Confocal immunofluorescent analysis of phospho-TSC2 antibody with HeLa cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used as a nuclear counterstain (blue).



Dot Blot

Image 2. Dot blot analysis of phospho-TSC2 antibody. 50ng of phos-peptide or nonphos-peptide per dot were spotted.



Immunofluorescence

Image 3. Confocal immunofluorescent analysis of phospho-TSC2 antibody with HeLa cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used as a nuclear counterstain (blue).